

This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + Refrain from automated querying Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at http://books.google.com/

OFFICIAL FORMULE



V125 TR3 1885 ILAIN D



LEVII COOTOPER! LANIE! FUIND





ILA NE




LEVII COOTPER LANE FUIND









-OF-

AMERICAN HOSPITI

COLLECTED AND ARRANGED

C. F. TAYLOR, M.D.,

EDITOR OF

THE MEDICAL WORLD.

Chpyright 1885, by C. E. Taylor

23

INTRODUCTION.

In collecting the matter for this book, all the principal, and nearly all the minor Hospitals in the United States were either visited or communicated with. Some hospitals have a printed list of formulæ; others have a few standard prescriptions, and still others have no established formulæ at all. The printed lists, the established prescriptions from those hospitals not having printed lists, and all other information of value, have been collected and are here presented. Omitting the lists of two or three hospitals that replied strangely "Not for publication," the writer feels that this collection is quite as complete as can well be made.

It is scarcely necessary to state that these formulæ do not represent the entirety of hospital practice. Certain mixtures being found useful in a large number of cases. they are prepared in large quantities to facilitate dispensing. This is important where a large number of cases are treated daily. Special cases, or cases presenting unusual indications, are treated according to indication presented. The writer has no admiration for doctors who blindly follow printed formulæ, and it is farthest from his desire that this book should encourage this large class of practitioners. The benefit intended is to place before the profession the remedies and combinations of remedies found most useful in our hospitals. The study of these cannot fail to give many hints and suggestions useful in private practice. It is hoped that the unlocking of these store-houses of experience will be considered a service to the profession.

C. F. TAYLOR.

PHILADELPHIA, August, 1885

CONTENTS.

Introduction Pharmacopoeia of the Hospital of the University of Penn-	3
sylvania (Philadelphia, Pa)	5
the Protestant Episcopal Church, Philadelphia, Pa Pharmacopæia of the Philadelphia Hospital, Philadelphia,	24
Pa	47
Prescribing Manual of the Children's Hospital of Philadel-	
phia, Pa Pennsylvania Hospital, (Philadelphia, Pa)	59 103
lefferson Collège Hospital, (Philadelphia, Pa.)	107
Formulary of the New York Hospital, (New York City) Roosevelt Hospital Formulary, (New York City)	134
Department of Public Charities and Correction, New York	
Bellevue Hospital	159
" Out Door Department Charity Hospital, Blackwell's Island	161
Insane Asylum, " "	174
Ninety-ninth Street Reception Hospital Hart's Island Hospital	175 176
Hart's Island Hospital	177
New York	179
Medical Formulary of the City Hospital of the City of Bos-	192
Chicago Marine Hospital. (Chicago, Ill.)	206
Cook County Hospital, (Chicago, Ill.)	22 I 22 3
Maryland Woman's Hospital, (Baltimore, Md)	225
St. John's Hospital, (Brooklyn, N. Y.)	226 228
Cincinnati Hospital, (Cincinnati, 0,)	230
Pa.)	. 232
U. S. Naval Hospital, (Washington, D. C.)	234 235
Home of Inebriates, (San Francisco, Cal.)	236

PHARMACOPŒIA

OF THE

Hospital of the University of Pennsylvania.

COLLYRIA.

Ŗ	Atropiæ sulphatisAquæ destillatæ	gr. iv f 3 j
!	М.	
R	Duboisiæ sulphatisAquæ destillatæ	gr. } f 3 j
1	М.	
R	EserinæAquæ destillatæ	gr. ij f z j
1	м.	
B	Zinci sulphatis	gr. iv f 3 j
1	M	
Ŗ.	Aluminis	gr. ij f 3 j
1	M. ·	•

OFFICIAL FORMULÆ

Ŗ.	Acidi galliciGlycerinæ	3 i
1	M. .	•
Ŗ.	Acidi tanniciGlycerinæ	3 ij
1	vi. -,	•
Ŗ.	Pilocarpiæ nitratis	gr. ij f Z j
1	M.	

GARGARISMATA.

Gargarisma Composita.

Ŗ	Potassii chloratis Rhois glabri	3 ij 3 j
M 3⁄3 P	I.—Sig. Put in a pint of boiling water a pint in an earthen vessel, and then strain	nd simmer to

GLYCERITA.

Giyeeritum Plumbi Subacetatis.

Ŗ	Plumbi acetatis	Э v 3 j
	Glycerinæ	3 vij

Heat for half an hour in a boiling glycerine bath, constantly stirring, and filter in a heated condition.

LINIMENTA.

1.—Linimoatum Tigli	i.
R Ol. tiglii	
М.	
2.—Linimontum Chloroformi C	
R Chloroformi	f z ij f z ss f z ss f z v
Tr. aconiti radicis	f g ss
Aquæ ammoniæ	fā ss
Ol. olivæ	fg v
М.	
3.—Linimentum Iodinii Com	positum.
R Chloroformi	fg ij fg vi
Tr. iodinii	f ž vj
M .	•
LIQUORES.	
1.—Liquor Picis Aikati	nus.
R. Picis liquidæ	fg ij g j fg v
Potassæ	. 3 j
Aquæ	•
M.—Sig. One teaspoonful to four or efuls of water; for external use.	eight tablespoon-
2.—Liquor Chi era lis	B
R Chloralis hydratis	gr. lxxx
Aquæ	fЗj
M.—Dose, one to two teaspoonfuls.	- · · .

LOTIONES.

1.-Lotio Nigra.

Ŗ.	Hydrargyri chloridi mitis Liq. calcis	gr. iij f Z j
N	vit	

2.—Lotio Pinmbi Acetatis et Opii.

Ŗ.	Plumbi acetatis	gr.	
	Tr. opii		j ss
	Aquæ	f 3	vj ss
1	л.		

MISTURÆ.

1.—Mistura Forri Arsonicolis.

Ŗ	Vini ferri	
	Liq. potassii arsenitis Syrupi	f 3 ss f 3 j ss
	Aquæ	f z iij
1	M.—Dose, one teaspoonful, diluted, aft	er meals.

2.—Mistura Acida Astringons.

Ŗ.	Acidi sulphurici aromatici Ext. hematoxyli	m	xl i
	Tr. opii camphoratæ,	f 3 f 3	iv iv,
N	1.—Dose, two teaspoonfuls.	- 3	

3.—Micture Crots Com	po.
R Mist cretze	
M.—Dose, a teaspoonful.	
4.—Misture Clei Phose	hors
R Ol. phosphorati	. 1
M.—Dose, one to two teaspoonfuls	•
5.—Mistura Potassii Chierai	le et f
R Potassii chloratis	. 13
M.—Dose, one teaspoonful.	•
S.—Misture Ferri Acc	4-4-
R Tr. ferri chloridi	· 7.

8.—Mistura Ferri Aperie	ns.
R Ferri sulphatis	gr. ij 3 ij fž j
9.—Mistura Potassii Iodidi F	ortior.
R Potassii iodidi	f j ss f j j of iodide
10.—Mistura Potassii lodi	di
R Potassii Iodidi	gr. xl f3 j
11.— Mistura Potassii lodidi Cor	nposita.
R Potassii iodidi	gr. xx gr. 1-6 f3 j
12.—Mistura Hydrargyri Com	nosita.
& Hydrargyri chloridi corrosivi	gr, ½ gr. lxxx

OF AMERICAN HOSPITALS.

	•	
	13.—Mistura Giberti.	
B.	Hydrargyri iodidi rubri Potassii iodidi	gr. ½ gr. xvj fg j
	Syrupi simplicis	fg vij
	M.—Dose, a tablespoonful.	
	14.—Mistura Hydrargyri et i	Ferri.
D	Hydrargyri chloridi corrosivi	
Ω×	Tr. ferri chloridi	gr. iv f % i
		13 J
	M.—Dose, ten drops.	
	15.—Misture Potassii Bron	aldi .
ĸ	Potassii bromidi 3	ij, gr. xl
	Aquæ cinnamomi	f ʒ j
	M.—Dose, one to two teaspoonfuls.	
	16Mistura Sedii Compes	site.
R	Sodii bicarbonatis	gr. xx
	Acidi carbolici	gtt. ij
	Acaciæ	B 1)
	Sacchariāā	q. s.
	Spts. lavandulæ comp	fg ij
	Aquæ	fzvi
	M.—Dose, a teaspoonful two hours after	<i>U ,</i>
		meais.
	17.—Mistura Oloi Morrhua Cer	nposita'.
R		ſ ž j
	Tr. iodinii comp	11 viii
	M.—Dose, one to four teaspoonfuls.	
	ar.— 2000, one to tour teaspoonius.	

	18. – Mistura Olei Morrhum et	Ætheris.	
R	Mist. ol. morrhuæ	f Z j	
	Ætheris	M ×vj	
]	M.—Dose, one to four teaspoonfuls.	•	
18	.—Mistura Olei Merrhuæ et Ca Phesphatis.	icis Lact	•
Ŗ	Ol. morrhuæ	f g ij	
]	Fiat emulsio et adde—		
	Syr. calcis lacto-phosphatis	f 3 ss	
_	20.—Mistura Hyescyami et M		
R	Morphiæ acetatis	gr. 1/4 f 3 j ss f 3 ij ss f 3 ss	
	Tr. hyoscyamiSyr. tolutanæ	fgjss fgijss	
	Aquæ	f ž ss	
1	M.—Dose, two teaspoonfuls.	3	
	21.—Misture Ammonii Chic	oridi.	
Ŗ.	Ammonii chloridi	gr. xx	
	Syr. scillæ	fg ij	
	Mist. glycyrrhizæ comp	fg vj	
]	M.—Dose, two to four teaspoonfuls.		
	22.—Mistura Cincheniæ A	cida.	
\mathbf{R}	Cinchoniæ sulphatis	gr. xii	
•	Acidi nitromuriatici dil	fg ss	
	Aquæ cinnamomi	f g v	
1	Ft. sol. Dose, two teaspoonfuls in water	, befo	

TERICAN TOOP J ZZ - Commercial C Priv come manaca Ponssi ogidi Vini colehici ceninis. ³cto-A THE CHARLES CHARLES THE CHARLES CHAR Syring M_Dose, a tablespoonful. Acidi nitromariatici dil....... Inf. gentiane comp. M.—Dose, two teaspoonfula, Liq. potaunii amenitia..... II (Prosesso **Sухар** M.—Done, one tempooned, dileger. meals. Spr. ather comp. Tr. inhelia Aque camphone **X**_n.

OFFIC	IAL	FOR	MUL	Æ

	28.—Mistura Doweesi	i.
	Tr. guaiaci comp	
	29.—Mistura Olei Ricin	ıl.
B	Ol. ricini	f 3 j gtt. iij
,	OO Michael Olei Membum Co	
R	30.—Mistura Oloi Morrhum Go Ol. morrhum Extract. malti	g ss
	31.—Mistura Croten-Chie	raiis.
	Croton-chloralis hydratis	gr. x f 3 ij f 3 vj spoonful, every
	PILULÆ.	
B.	1.—Pliula Opii ot Plumb Pulv. opii Plumbi acetatis	gr. 1/3 gr. ij
_	• •	

2	SPITALS.
D., Pilula On	il et Acidi Tann
Acid: opii	Of Acidi To-
Pulv. opii	anni
et ft. pil. No. 1.	9

3.—Pilula	9
vdra- Hydra	Manual Marie Control
3. Pilula Hydras ydrargyri iodidi viridi nfectionis rosæ ft. pil. N	Jor lodidi Vi.
A TOSE TOSE	*****
ft. pil. No. 1.	gr.
	gr.
-Pilula Col-	
colocuration	histo o
Pliula Colocyaticolocynthidis comp	Composit:
colocynthidis comp pelladonnæ	*********
rui	
pil. No. 1.	87. 1/s
740. I.	gr. j

	···· 1.	
3.—PI B. H	iula Hydran	-
Confectio	ri iodidi viridi nis rosæ No. 1.	lyri lodidi Vi
-	-10. I.	*****
R Ext. color	la Colocyuth	
Ext. bellado	a Colocyata nthidis comp nnæ	Composi
M. et ft. pil. N	nnæ.	gr. ij
	o, 1.	gr. j
R. Resinæ podort		0
Pulv. rhei Ext. hall	Podophytti ylli	posita.

M. et ft. pil. No. 1.

	7.—Pilula Arsenicelis Com	posita.
R.	~	gr. 1-35 gr. j ss gr. ij
	•	
	8.—Pliuia Cinchoniæ cum	Ferro.
Ŗ	Cinchoniæ sulphatis Pulv. ferri Strychniæ M. et ft. pil. No. 1.	gr. ij gr. j gr. 1-30
	9.—Pilula Argenti Nitrat	is.
Ŗ.	Argenti nitratis	gr. ¼ gr. 1-12
1	M. et st. pil. No. 1.	
	10.—Piiula Ferri Cempos	ita.
R N	Acidi arseniosi	gr. 1-35 gr. 1-35 gr. 1-5 gr. j ss gr. ij ss
_	11.—Pilula Arsenicalis.	1
B.	Acidi arseniosi	gr. 1-20 gr. ij
		4.5

	12Pliula Auri et Sodii Chioridi. Auri et sodii chloridi
	13.—Piluia Gamphora Monobromidi. Camphora monobromidi
	PULVERES.
-	1.—Pulvis Bismuthi Gempositus. Bismuthi subnitratis
Ŗ.	2.—Pulvis Soda Gompositus. Sodii bicarbonatis
	3.—Pulvis Bismuthi. Bismuthi subnitratis gr. xv ct. chart. No. 1.

	4.—Pulvis Santonini.
\mathbf{R}	Santonini gr. 1/2 Sacchari gr. ij
	Saccharigr. ij
F	t. chart. No. 1.
	5.—Pulvis Rhei Compositus.
R	Pulv. rhei gr. iii
•	Sodit bicarbonatis gr. x
	Pulv. zingiberis gr. ij
M	I. et ft. chart. No. 1.
	9.—Pulvis Buchu Compositus.
R.	
	Buchu
	Folii belladonnæ gr. viij
wate	I. et ft. chart. No. 1. Sig. Put in a pint of boiling er, and when cold, strain; dose, a wineglassful three es a day.
	7.—Pulvis Gentianæ et Quassiæ.
R.	Gentianæ
12	Quassiæ
	Činnamomiāā ʒ ij
M	I.—Sig. Put in a pint of boiling water, and when
	, strain; dose, a wineglassful three times a day.
	8.—Pulvis Juniperi Compositus.
\mathbf{R}	Juniperi
	Potassii bitartratis
wat	 et st. chart. No. 1.—Sig. Put in a pint of boiling er, and when cold, strain; dose, a wineglassful three es a day, before meals.

9.—Pulvis Anthemidis et Quassiæ.
R Anthemidis
M. et ft. chart. No. 1.—Sig. Put in a pint of boiling
water, and when cold, strain; dose, a wineglassful three
times a day, before meals.
10.—Puivis Calumbæ Compesitus.
R. Calumbæ
Zingiberis āā 3 ss Sennæ 3 ij
M. et ft. chart. No. 1.—Sig. Put in a pint of boiling
water, and when cold, strain; dose, a wineglassful three times a day.
44 Puluis Olmiskium Comassikus
11.—Pulvis Cimicifugæ Compositus.
R. Cimicifugæ
Zingiberis
M. et ft. chart. No. 1Sig. Put in a pint of boiling
water, and when cold, strain; dose, a wineglassful three times a day.
12.—Pulvis Petassii Bromidi Cempositus.
R Potassii bromidi
Ferri sulphatis exsiccatæ
Zingiberis
M. et st. chart. No. 1.—Sig. Put in a pint of boiling
water, and when cold, strain; dose, a tablespoonful three times a day.

	13.—Pulvis Hydrargyri et	Sodii.
Ŗ	Hydrargyri chloridi mitis Sodii bicarbonatis	gr. ½ gr. x
I	t. chart. No. 1.	•
	14.—Pulvis Cincheniæ et Po	opsinæ.
R	Cinchoniæ sulphatis	gr. iij
N	1. et ft. chart. No. 1.	
	15.—Pulvis Ammonii Chic	oridi.
Ŗ.	Ammonii chloridiāā	3 ij
dos	1. et st. chart. No. 1.—Sig. Put in ½ e, a tablespoonful four times a day.	
	19.—Pulvis Cinchonidiæ Con	npositus.
Ŗ	Cinchonidiæ sulphatis Extr. glycyrrhizæ Sacchari	gr. iij
N	I. et st. chart. No. 1.	8
•	0	
	SUPPOSITORIA.	
	1.—Suppositorium Morp	hiæ.
Ŗ.	Morphiæ sulphatisOl. theobromæ	gr. ss q. s .
I	t. suppos. No. 1.	-

2.—Suppesiterium lodoformi.			
R	IodoformiOl. theobromæFt. suppos. No. 1.	gr. v q. s.	
3.		eliadonnæ.	
Ŗ	Morphiæ sulphatis	gr. ss gr. j ss q s.	
	 LSuppositorium Morphiæ A	stringens.	
B	· · · · · · · · · · · · · · · · · ·	gr. ss gr. j ss gr. v q. s.	
5	. –Suppositorium Morphiæ Co	mpesitum.	
₽.	Sodii boratis	gr. v gr. ss gr. j ss q. s.	
	6.—Suppositorium Aeidi Ta	nnici.	
₽.	Acidi tanniciOl. theobromæ	gr. v q. s.	

TINCTURÆ.

1.—Tinctura iedinii et Opii.				
R Tr. iodinii				
2.—Tinctura Saponis Viridia.				
R Saponis viridis				
UNGUENTA.				
1.—Unguentum Diachyli.				
R Plumbi oxidi				
2.—Unguentum Belladonnæ Compositum.				
R. Ext. belladonnæ				

	3.—Unguentum Hydrargyri Ca Ungt. hydrargyri	•	situm ij
	M.	ວ	•)
	4.—Unguestum Veratri	æ .	
Ŗ	Veratriæ Adipis	D S	ij j
]	M. et ft. ungt.		
1	5.—Unguentum Hydrargyri Ox	idi I	Flavi.
Ŗ	Hydrargyri oxidi flaviUngt. petrolei	gr. 3	xvj j
1	М.		
	6.—Unguentum Hydrargyri An	nme	niati.
Ŗ			lxxx
1	М.	3	•

PRESCRIBING MANUAL

OF THE

DISPENSARY

OF THE

Hospital of the Protestant Episcopal Church

OF

PHILADELPHIA.

In prescribing, the Physician will write the number of the prescription, its title, the quantity to be dispensed, and the directions for using, e. g:—

For.....

R. No 2.—Liquor Potassii Bromidi, 13 iij Sig One teaspoonful, diluted, three times a day. Date. Signature.

Unless otherwise directed, the apothecary will always dispense three fluid ounces (f 3 iij) of the fluid preparations, when the dose ordered is one fluid drachm (f 3 i), and six fluid ounces (f 3 vi) when the dose is half a fluid oune (f 3 ss).

-LAICAN HOSPITALS

MEDICINES

INTERNAL ADMINISTRAT

Note. The following formulæ are desi adults only.

1.—Liquor Poi	LIQUORES.
	LIQUORES. Passil Browld (Morveus tive.)
M.—Dose, 13 j-	gr. v

M.—Dose, f 3 j-iv, diluted, three times daily.

MISTURÆ.

- Misturo Arsonicalis (Alterativo). R. Lig. potassii arsenitis.....
 - Infus. gentianæ comp..... q. s...ad.
- M.—Dose, [3], diluted, thrice daily, after meals.
- 3. Mistera Hydrargyri et Fami o R Hydrargyri chlorid:

5.—Mistura Petassii lodid	li (b).
Potassii iodidi	gr v m x m xx f g j aily.
6.—Mistura Creasoti (Anta	icid).
R. Aquæ creasoti	f 3 j gr. x f 3 ss
7.—Mistura Sodii Bicarbena	tis (a) .
R Sodii bicarbonatis	gr. x f 3 j f 3 ss
8.—Mistura Sodii Bicarbona	tis (b).
B. Sodii bicarbonatis	gr. v m xx fg j
9.—Mistura Acidi Salicylici (An	tipyretic).
R. Acidi salicylici	gr. v m xl m v m xv
M.—Dose, f 3j, every two or three hou	rs.

10.—Mistura Lobelia Comp. (Antispasmodic)				
R	Tinct. lobeliæ	m x m xxx m x fg j		
11.	.—Mistura Acidi Sulphurici (a—	Astringont)		
R	Acidi sulphurici aromat	m x m v f3 j f3 ij		
	12.—Mistura Cretæ Com	p.		
	Tr. opii	m v fgj m xv fgss		
13.—Mistura Pepsinæ Cemp. (Digestant.)				
3.	Pepsinœ	gr. x m xv f 3 ss f 3 ij		

	14.—Mistura Digitalis Comp. (I	Diuretic).
B	Tinct. digitalis	m, v
	Potassii acetatis	gr. xx
	Spt. ætheris nitrosi	í g j
	Aquæq. sad.	f 🖁 ss
	M.—Dose, f 3 ss, three times a day.	_
15	5.—Mistura Ammon. Cemp. (Ex	(pectorant).
ĸ	Copaibæ	m v
	Ext. glycyrrhizæ pulv	gr. x
	Ol. anisi	m ij
	Misturæ ammoniaciq. sd.	fg j
	M.—Dose, f 3j, every three or four hou	rs.
	16.—Mistura Ammonil Chlor	idi (a).
R		gr. v
	Mist. glycyrrhizæ comp	ſЗj
	M.—Dose, f 3 j-ij, every two or three ho	ours.
	17.—Mistura Ammonii Chier	idi (b).
Ŗ		gr. v
	Potassii iodidi	gr. ij
	Glycerinæ	m xx
	Syr. senegæ	m xv
	Aquæq. sad.	fзj
	M.—Dose, fzj, every two or three hour	's,
	18.—Mistura Petassii Cya	nidi.
R		_
	Morphiæ sulph	gr. 1-16
	Syr. tolutani	f 3 ss
	Aquæq. sad.	fЗj
	Dose, f 3 j, every six hours.	

19	.—Mistura Ergotæ Comp. $(oldsymbol{a}$ —Ha	emestatic).
Ŗ	Ex. ergotæ fluidi	m, xx
	Tinct. digitalis	m ijss
	Tinct. opii deodorat	m iij
_	Aquæ cinnamomiq. sad.	fзj
1	M.—Dose, f 3j, every one, two or three	hours.
	20.—Mistura Ergotæ Com	p. (<i>b</i>).
Ŗ	Ex. crgotæ fluidi	fg ss
	Potassii bromidi	gr. xv
	Aquæ cinnamomiq. sad.	f g j
]	M.—Dose, f z j-ij, p. r. n.	
•	21.—Mistura Ferri Acida (Laxe	ative end
•	Cathartic).	
R		gr. j
	Magnesii sulphatis	gr. xl
	Acidi sulphurici dil	m vij
	Infus. quassiæq. sad.	f 🖁 ss
j	M. —Dose, f 3 ss, once or twice daily, w	ell dilut ed.
	22.—Mistura Sodii Bicarbona	atis $(oldsymbol{c})$.
\mathbf{R}		
	Aloes socotrinæāā	gr. x
	Spirit. lavandulæ comp	f 3 ss f 3 ss
_	Aquæq. sad.	1 3 ss
1	M.—Dose, f z ij-iv, once or twice daily.	
23	.—Mistura Ammonii Carbenati	s (a—Stim-
	ulant.)	- (13 Guill
Ŗ.	Ammon. carbonatis	gr. v
	Syr. tolutani	f 3 s8
	Mucilag. acaciæ	fĝj
1	M.—Dose, f 3 j-ij, every three hours.	

2	24.—Mistura Nitro-Muriatici (1 Antiperiodic).	romic and
\mathbb{R}	Acidi nitro muriatici dil	m x
	Infus. gentianæ compq, sad.	тх fgj
	M.—Dose, f3j, diluted, three times da	ily.
	25.—Mistura Cinchonid	iæ.
Ŗ	Cinchonidiæ sulphatisAcidi sulphurici dil	gr. ij m iv føj
]	M.—Dose, f 3 j-ij.	.9 J
	26.—Mistura Ferri Aceta	
Ŗ	Liquor ammonii acetatis	fg j, m xl
	Tinct. ferri chloridi	m x
1	M.—Dose, f 3 j-ij, three or four times a	day.
	* ***********************************	
	27.—Mist. Forri Acetat. et Str	ychniæ.
R	Strychniæ sulphatis	gr. 1-30
	Mist. ferri acetatis	fg ij
1	M.—Dose, f z j-ij, thrice daily.	
	28.—Mistura Ferri Com	p.
R.	Strychniæ sulphatis	gr. 1-60
	Cinchoniæ sulphatis	gr. ij
	Tinct. ferri chloridi	mχ
	Acidi muriatici dil	mγ
_	Aquæq. sad.	fg j
1	M.—Dose, f 3 j, diluted, thrice daily	

	29.—Mistura Gentianzo Co	emp. \
R .	Ferri et ammonii citratis	gr. x f3 j f3 j f3 j f3 ss
	 30Mistura Strychniæ ((a).
Ŗ.	Strychniæ sulph	gr. 1-48 m x m xx f 3 ij
1	M.—Dose, f 3 ij, diluted, thrice daily.	•
	31.—Mistura Strychniæ (b).
Ŗ.	Strychniæ sulphatis	gr. 1-48 m v

hours.

33.—Pilula Quinim et Digitalis (Antipyretio).
R. Quiniæ sulphatis gr. ij Pulv. digitalis gr. ss
Pulv. opii gr. ¼
M. et ft. pil. no. j.—Dose, one pill every six hours.
22. 3. 1. p. 2. 2. 3. 2. 3. 5. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.
34.—Pliula Argenti Nitratis (Astringent).
R Argenti nitratis gr. 1-6
Ext. gentiannæ gr j
M. et ft. pil. no. j.—Dose, one pill every four or six
hours.
35.—Piluia Argenti Nitrat. et Opii.
R Argenti nitratis
Pulv. opii
Ext. gentianæ gr. j
M. et ft. pil. no. j.—Dose, one pill every four or six
hours.
ulaudia v
36.—Pilula Cupri Suiphatis.
R Cupri sulphatis gr. 1-6
Pulv. opii gr. 1-6
Ext. gentianæ gr. j
M. et ft. pil. no. j.—Dose, one pill every four or six
have

37.—Pilula Celccynth Cemp. (Cathartic).
& Ex. colocynth comp gr. ij
Res. podophyllin gr. 1/8
Ex. belladonnæ gr. 1-6
Ex. gentianæ gr. j
Olei carui gtt. ss
M. et ft. pil. no. j.—Dose, one pill at night, or night and morning.
——————————————————————————————————————
PULVERES.
38.—Pulvis Sadii Comp. (Antacid).
R Sodii bicarbonatis gr. v Bismuthi subcarbonatis gr. x
M. et st. chart. no. j.—Dose, one powder thrice daily.
39.—Pulvis Kameeim (Anthelmintic).
R Pulv. kameelæ 3 j
Ft. chart. no. j.—Dose, one or two at night, or night
and morning.
40.—Pulvis Bismuthi (Astringent'.
R Bismuthi subnitratis 3 j
Ft. chart. no. j.—Dose, one powder thrice daily.
41.—Puivis Bismuthi et Ipecac.
R Bismuthi subnitratis gr. x
Pulv. ipecac. comp gr. iij
M. et st. chart. no. j.—Dose, one powder every three
or four hours.

	42.—Pulvis Bismuthi et Merphiæ.
Ŗ.	Bismuthi subnitratis gr. x Morphiæ sulphatis gr. ½
	1. et ft. chart. no. j —Dose, one every four or six
hou	rs.
43	.—Pulvis Bismuthi et Pepsinæ (Digestant)
R	Bismuthi subnitratis
~~	Pepsinæ gr. x
N	1. et ft. chart. no. j.—Dose, one powder thrice daily.
	44 Pulule Redii et Penelum
	44.—Pulvis Sedii et Pepsinæ.
Ŗ	Sodii bicarbonatis gr. v
_	Pepsinægr. x
N	1. et ft. chart. no. j.—Dose, one powder thrice daily.
	 .
	45.—Pulvis Buchu Cemp. (Diuretic).
R.	Foliæ buchu
*	Folise uvse ursi
	Sem, lini
Ŋ	I. et ft. chart, no. j.—Sig. Add two pints of boiling
wat dai	er; when cold, strain. Dose, a wineglassful thrice
441	17.

MEDICINES

I'OR

CHILDREN.

In the following formulæ the doses have been proportioned for children who have reached the age of two years:

MISTURÆ.

•	16.—Mistura Olei Merrhuæ (Al	terative).
B	Olei morrhuæSyr.calcii lacto-phosphat	fg ss
	Aquae calcis	m xv
ľ	M.—Dose, f z j-ij, three times daily	
	47 Mistura Potassii Arse	nitis.
R	Liquor potassii arsenitis	ηjij
	Syr. simplicis	f 3 ss
	Aquæq. sad.	ſĝj
ľ	1Dose, f z j, thrice daily.	
	48.—Mistura Potassii iodid	ii (c).
Ŗ.	Potassii iodidi	gr. j
	Syr. sarsap. comp	
	Aquæāā	fg se
1	MDose, f g j-ij, thrice daily.	

49.—Mistura Antacida (Antacid).		
	Liquor calcis	f 3 ss
	50. – Mistura Sodii et Rh	ei.
R M	Sodii bicarbonatis	gr. ij m xv f z j
51.	-Mistura Chioral. Hydret.(Anti:	epasmodic)
Ŗ	- ,	gr. ij gr. v f z ss
	52.—Mistura Belladonna	10.
R.	Tinct. belladonnæ	m ij gr. v f 3 ss f 3 j
		•
53	.—Mistura Acidi Sulphurici ($m{b}$ —	Astringent)
B.	Acidi sulphurici dil	m ij m viij m v fg j rs.

54.—Mistura Bismuthi.

R.	Bismuthi subnitratis	gr. vj m j m xx f g j rs.
	55.—Mistura Antifehrijis (Dia _l	pheretic).
R	Potassii citratis	gr. ij
	Syr. limonis	m, x
	Liquor ammon. acetatq. sad	făi
3	M.—Dose, fgj, every two or three hour	rs.
	56Mistura Pepsina (Dige	stant).
Ŗ.	Pepsinæ saccharat	gr. iij m iij m v f3 j
,	I.—Dose, f 3 j, three times daily.	•
	T Michiga Defensii Asstatis (D 1

57.—Mistura Potassii Acetatis (Diuretic.)

Ŗ	Potassii acetatis	gr. ij ss
	Sp. juniperis comp	m, v
	Sp. ætheris nitrosi	m x
	Syr. simplicis	m, xx
	Aquæq. sad.	fǯj
7	I -Dose, f 3 j, every three or four hou	rs.

	58Mistura Ammor. Carbo	nat. (<i>b</i>).
Ŗ	Ammon. carbonatis	gr. j
	Aquæ	m, xx
	MDose, one teaspoonful every two o	r three hours.
5	9Mist. Ammonii Chloridi ($oldsymbol{c}$ -E	xpectorant)
R	Syr. ipecacuanhæ	m, v
•	Ammonii chloridi	gr. ij
	Syr. tolutan	g ss
	Aquæq. s	f <mark>g</mark> j
	M.—Dose, f 3 j, every three hours.	0,
	22. 2 300, 1 3), 0 100, 11120 11011101	
		
	60.—Mistura Ipecac. Co	mp.
R	Vini ipecacuanhæ	m, iij
7.	Sp. ætheris nitrosi	• ,
	Tr. opii camphorat	m iv
	Syrupi tolutani	mχ
	Liq. ammonii acetatisq sad.	ſg j
	M.—Dose, f z j, every three hours.	
6	I.—Mistura Aloes et Ferri (La Cathartic).	xative and
Ħ	Tinct. aloes et ferri	m x gr. 1-6 f3 j

62.—Mistura Magnesii et Opii.

R	Magnesii sulphatis	gr. v
	Tr. opii deodorat	m, ss
	Syrupi	m. x
	Aq. menth. pipq, sad.	fʒj
R	1.—Dose, f z i, every three hours.	_

63.—Mistura Ferri et Cinchonidiæ (Tonic and Antiperiodic).

Ŗ	Cinchonidiæ sulphatis	gr. j m. ij
	Syr. zingiberis	f g ss
	Aquæq. sad.	ſā.j

84.-- Mistura Potassii Chiorat. (a).

Ŗ	Potassii chloratis	gr. ij	SS
	Tinct. ferri chloridi	m, ij	SS
	Ac. muriatici dil	m, ij	
	Syr. zingiberis	fzss	,
	Aquæq. sad.	fzi	

M.—Dose, fzj, every two or three hours.

65.-- Mistura Potassil Chlorat. (b).

Ŗ	Potassii chloratis	gr. ij M. ij
	Glycerinæ	m x
	Aquæq. sad.	f g j

M.—Dose, f z j, three to five times daily, in water.

PULVERES.

66.—Pulvis Hydrarg. et Sodii ()	Antacid).
R Hydrarg, chloridi mit	gr. ij
M. et ft. chart. no j.—Dose, one every	2 or 3 hours.
67.—Pulvis Sodii et Bismu	thi.
R Sodii bicarbonatis	
Bismuthi subcarbonatis	gr. ij
M. et ft. chart. no. j.—Dose, one powde	er thrice daily.
68.—Pulvis Santonini (Anthel	mintic).
R Santonini	gr. ss
Sacch. albi	gr. j ss
M. ef ft. chart. no. j.—Dose, one or morning and evening, the second to be follow	two powders
morning and evening, the second to be follow	red by a purge.
69.—Pulvis Bismuthi et Opii (As	tringent).
R Bismuthi subnitratis	gr. v
Pulv. opii	gr. 1-12
M. et st. chart no. j Dose, one powder	every 3 hours.
70.—Pulvis Cretæ Præparatæ	Comp.
R Cretæ præparatæ	gr. iv
Pulv. ipecac co	gr. ss
Puly, aromat	
M. et ft. chart. no. j.—Dose, one powde	r every two or
three hours.	
	The same of the same of
71.—Pulvis Hydrarg. Chloridi Mit.	(Laxative)
71.—Pulvis Hydrarg. Chloridi Mit.	(Laxative)
	gr. j
71.—Pulvis Hydrarg. Chloridi Mit. R Hydrarg. chloridi mitis	gr. j gr. ;

PREPARATIONS

FOR

EXTERNAL USE.

The various fluid preparations embraced in this list are to be kept and dispensed in colored and roughened bottles, labelled "For External Use."

GARGARISMATA.

72	2.—Bargarisma Potass. Chlore Carboliel.	at. e	t Acidl
R	Potassii chloratis	gr.	xx
	Acidi carbolici	gr.	
	Glycerinæ		lxxx
	Aquæ q. s a d.	f 3	J
	M.—Sig. Add, f 3 ss to f 3 iv of water.		•
73		t. et	Forri.
R	Potass. chloratis	gr.	xx
	Tinct. ferri chloridi	m f 3	х×
	Aquæq. sad. M.—Sig. Add f 3 ss to f 3 iv of water.	13	,J
	74.—Bargarisma Sedil Bibo	orati:	8.
R	Sodii biboratis	gr.	XΨ
	Glycerinæ	f 3	ij
	Aquæq. sad.	fξ	j
	м.	Ŭ	-

GLYCERITA.

	75.—Glyceritum Tr. Forri Cl	alori	dĺ.
Ŗ	Tinct. ferri chloridi		
	Glycerinæāā	f Z	SS
1	M.		
	70 Okronitum Indial		
_	76.—Glyceritum lodini	١.	
R	Liq. iodinii comp	fξ	
	Glycerinæūā	13	55
1	M 0		
	LINIMENTA.		
	77.—Linimentum Aconiti C		
R	Tinct. aconiti rad	ous.	•
ſκ	Chloroformi		
	Tr. opii		
	Olei olivæāā	f 3	j
	Lin. saponis camphq. sad.	f₹	j
1	M		
	78.—Linimontum Stimul	ans.	
R			ii
1,2	Lin. ammoniæ	f 3	iv
1	М.	·	
			
	LIQUORE S .		
	· 79.—Liquor Acidi Carbolic	i (a	١_
R		gr.	
174	Aquæ	f 3	· i
	M.	- 0	•

	30.—Liquer Acidi Carbolic	l (b).
R.	Acidi carbolici	
	Aquæ	gr.xx f3 j
1	M.	· ·
	81.—Liquor Argenti Nitratia	s (a).
Ŗ	Argenti nitratis	gr. ij fīži
	Aquæ destillat	fЗj
3	1 .	
		
	82.—Liquor Argonti Nitratis	B (b).
Ŗ.		•
-	Aquæ destillat	gr.x f37 j
1	vI.	•
. •		
	83.—Liquor Argenti Nitrati	s (<i>c</i>).
ß		• •
	Aquæ destillat	gr. xxx f Z j
7	vI.	0,
	·	
	84.—Liquor Atropiæ (a)-
R	Atropiæ sulphatis	gr. ss
	Aquæ destillat	gr.ss f3; j
1	M.	•
		
	85.—Liquor Atropiæ (b)_
R	Atropiæ sulphatis	gr. iv
	Aquæ destillat	fij
1	м.	

90.—Pulv. Plumbi. Acetat. ot Zinci.

Plumbi acetatis.....
Zinci acetatis.....
M.—Sig. Dissolve in a quart of water.

	91.—Pulvis Querc. Alb. et Zinci.
Ŗ	Querci albæ
	I.—Sig. Add a quart of boiling water, let stand for hours, and strain.
	 92.—Pulvis Quassiæ.
R	Quassiæ ligni 3 ij
S	ig. Add one pint of boiling water, let stand till cold, strain.
	o
	UNGUENTA.
	93.—Unguentum Diachyli.
R	Lithargyri
C	oque, dein adde
	Olei lavandulæ m iv
M	. et ft. ung.
	94.—Ung. Iodinii et Belladonnæ.
R M	Ung. iodin. comp

OFFICIAL FORMULÆ

	95.—Ung. lodof erm i.		
K	IodoformiAdipis	3	j j
	M. et ft. ung		
	G6.—Ung. Stramonii et Ac. T	'Ann	iel.
Ŗ	Ung. stramonii		
	M. et ft. ung.	_	

PHARMACOPŒIA

OF THE

PHILADELPHIA HOSPITAL.

CHARTÆ.

Charta Arsenicalis Composita.

Ŗ.	Belladonnæ foliæ	gr.	xcvj
	Hyoscyami foliæ		x lviij
	Extracti opii	gr. gr.	lxx x
F	iat infusum, et adde		
	Potassii nitratis Potassii arsenitis		clx cccx>
F	iat solutio, et –		
	Saturate bibulous paper and dry for u	se.	

GLYCERITA.

Glyceritæ Olei Ricini.

Ŗ.	Olei cinnamomi	gtt.	xxiv
	Glycerini	f₹	iv
N	Misce.		

INFUSA.

Infusum Sennæ Compesitum.

R Sennæ foliæ	z viij z iij O vj
Macera per horas duas, cola, et adde-	
Tincturæ sennæ compositi	O ij
Misce —Signa. Dose, a tablespoonful.	

LINIMENTA.

Linimentum Chlorofermi Compositum.

Ŗ	ChloroformiTincturæ aconiti radicis	ſ ℥	ij
	Aquæ ammoniæ	f 3	SS V
I	Fiat linimenta.	Ŭ	

	Linimentum Seponis Virido.
3 .	Saponis viridis
l ilt	Fiat solutio. Liquefac cum leni calore. Misce c ra.
1	Linimentum Terebiathinas Compositum.
B.	Olei terebinthinse
1	Flat linimenta.
	Liquores.
	Liquer Bremini.
Ŗ.	Bromini 3 j Aque f 3 ij Potassii bromidi q. s.
	Fiat solutio. Signa. For physicians' use, only.
L	iquer Plumbi Subscelatis cum Opii (Lead water and Laudanum).
Ŗ.	Tincture opii
	and Oil

Misce.

MISTURÆ.

Mistura Antirheumatica.

R	Potassæ nitratis	7 i
~~	Vini colchici radicis	3 j 13 j
	Spiritus ætheris nitrosi	fži
	Syrupi guiaci	f¥ ii
	Olei gaultheriæ	gtt. vj
	Aquæq. sad.	f ž vj

Misce.—Signa. Dose, a tablespoonful every two hours.

Mistura Arsenicalis Composita.

R.	Liquoris arsenici chloridi	f 3 ss f 3 j ss
-	Tincturæ ferri chloridi	f 🕇 j ss
	Cinchoniæ sulphatis	ž ij
	Strychniæ sulphatis	gr. ij
	Syrupi	
	Aquæq. sad.	f Z vj

Fiat mistura. Signa-Dose, a teaspoonful.

Mistura Astringens.

B.	Acidi sulphurici aromatici	f g ij
•	Extracti hæmotoxyli	Ž ij
	Tincturæ opii camphoratæ	f 3 ss f 3 vj
	Syrupi zingiberisq, sad.	f℥ vj

Misce secundem artem.—Signa. Dose, a tablespoonful.

Mi	stura Cosmetica (Goddard's Cosmetic Le- tion).
R 1	Tincturæ benzoini
	Mistura Cretæ Composita.
R	Tincturæ catechu
ful.	Isce secundem artem.—Signa. Dose, a tablespoon-
Mi	stura Ferri Chloridi Composita (Basham's Mixture).
R	Liquoris ammoniæ acetatis

Mistura Sodæ (Soda Mint).

Ŗ	Sodæ bicarbonatis	3 ij	i
	Spiritus ammoniæ aromatici	git. x	
	Aquæ menthæ viridis	Ϊ ∄ v	iij

Misce.—Signa. Dose, one or two tablespoonfuls three times a day.

Mistura Sodæ Composita.

Ŗ	Sodæ bicarbonatis	3 gtt.	
	Syrupi acaciæ	fZ	ij
	Spiritus lavandulæ compositæ Aquæq. sad.	f 3	

Fiat mistura. Signa.—Dose, tablespoonful two hours after meals.

Chlorodyne.

Ŗ	Chloroformi	f 3 ss f 5 j ss
	Olei menthæ piperiæ	gtt. viij
	Oleoresinæ capsici	gtt. ij
	Extracti cannabis Indicæ	gr. vj
	Morphiæ muriatis	gr. xvj
	Acidi hydrocyanici diluti	m lxv
	Acidi hydrochlorici diluti	fgj
	Glycerinæ	
	Méllis	fZ iv

Fiat mistura secundem artem. Signal.—Dose, 15 to ao drops.

MICAN HOSPITALS.

Mistura Zolilekofferi (Zellickoffer's Mixt

R		lidiāā	z ij ss
	Vini colchid	radicisamomi	f 3 jss
		q. sad.	Οj
	Fiat mistura.	Signa.—Dose, a tablespe	oonful

PILULÆ.

Pilula Alcos Composita.

Ŗ	Pulveris aloes socotrinæ Ferri sulphatis exsiccatæ		3	S S
	Terebinthinæ albæ	āā გ	3	ij
1	Misce, et fiant pilulæ cxx. S	ig na .— Each p	ill	contai

Misce, et fiant pilulæ cxx. Signa.—Each pill contain 2 grs. of aloes, and 1 gr. each of iron and

Pilulæ Cincheniæ ot Arsenici.

\mathbf{R}	Cinchoniæ sulphatis		
	Ferri redacti	3	SS
	Extracti nucis vomicæ	gr.	xxx
	Acidi arseniosi	gr.	νj

Misce, et fiant pilulæ cxx. Signa.—Each pill contains 1-20 gr. of arsenic, 1/2 gr. of nux vomica, and 2 grs. each of iron and cinchonia.

Piluiæ Colocynthæ cum Belladennæ.

ĸ	Extracti belladonnæ	gr.	χv
	Extracti colocynthidis compositæ	_	:::
	Pulveris aloes socotrinæāā Olei anisi	3 ett.	•
	Old mindiminiminiminiminiminiminiminiminimin	E.L.	~~

Misce, et fiant pilulæ cxx. Signa —Each pill contains 1 8 gr. of belladonna, and 1 ½ grs. each of aloes and colocynth.

Pilulæ Opii cum Piumbi Acotatæ.

Ŗ	Pulveris opii	gr. x	ĸl
	Plumbi acetatæ	3 s	iS.

Misce, et fiant pilulæ cxx. Signa.—Each pill contains ½ gr. of opium, and 2 grs. of acetate of lead.

Pilulæ Cinchonlæ Compositæ.

R	Cinchoniæ sulphatis	
-	Ferri redacti	₹ ss
	Extracti nucis vomicæ	gr. xxx

Misce, et fiant pilulæ cxx. Signa.—Each pill contains χ gr. of nux vomica, and 2 grs. each of iron and cinchonia.

	Pilula Podophylli Compo	sitas.
R.	Resinæ podophylli	
	Extracti colocynthidis composita	J
	Extracti hyoscyami	3 ij
I	Misce, et fiant pilulæ cxx. Signa.—Eac	h pill contains
	gr. of podophyllin, and I gr. each of	colocynth and
hyc	scyamus.	
	Piluia Rhei et Gentiana	B.
R		3 ss
	Extracti gentianæ	3
	Extracti hyoscyami	Зij
1	sisce, et fiant pilulæ cxx. Signa.—Eac	h pill contains
2 g	rs. of rhubarb, and I gr. each of gentia	n and hyoscy-
am	us.	
	PULVERES.	
	Pulvoris Glycyrrhize Comp	osita.
R.	Pulveris sennæ	
	Pulveris glycyrrhizæ radicisāā	₹ vj
	Pulveris fœniculi	
	Sulphuris loti	Z iij Z xviij
	Sacchari albæ	
MisceSigna Dose, a teaspoonful at bed time.		
_	Puiveris Sodæ Composi	
R	Bismuthi subnitratis	gr. v
	Sodæ bicarbonatis Pulveris zingiberi	
	Pulveris calumbæāā	gr. ijss
1	liera	8.4.1

Misce.

SYRUPI

B. Chloral hydrat
chloral.
Syrupus Gualaci.
Pulveris guaiaci resinæ
five grs. of guaiacum.
Syrupus Pectoralis.
Ammonise muriatis
Misce.—Signa. Dose, a dessertspoonful.
Syrupus Potassii lodidi.
Potassii iodidi

Syrupus Potassii Iodidi Composita.

TINCTURÆ.

Tincture Arometics.

Coriandri fructus	7 ij 7 ii \$s
Glyceringe	f∄ v f∄ vi
Alcoholis diluti, q. s., ut fiant tinc-	Oii
	Angelicæ fructus

Signa.—A pleasant vehicle for administering nauseous remedies.

Tinctura Ferri Composita.

R	Cinchoniæ sulphatis	3 j
	Strychniæ sulphatis	
	Tincturæ ferri chloridi	gr. ij f 🕇 j
	Syrupi	0.
	Aquæ	f Z v iij
1	Missa sasumdam antom. Sisma Dos	a a taganaan

Misce secundem artem. Signa.—Dose, a teaspoonful three times a day.

	Tinctura Saponis Viride cu	m P	iels.
Ŗ	Saponis viridis		
	Spiritus methlyciឆឺធឺ	3	j
	Misce cum leni calore.		
	Tinctura Styptica.		
R		7	i
·	Saponis	13 13	j ij iv
	Fiat mistura secundem artem.	Ü	
	UNGUENTA.		
	Unguentum Piumbi Oxi	idi.	
Ŗ	Emplastræ plumbi	f 3	j ij
	Misce cum leni calore.		
	Unguentum Zinci Oxidi Beni	zoat	um.
R	_	3	
~	Tincturæ benzoini	gtt.	xl
	Adipis	_ Z	vij
	Fiat unguentum.	Ü	•

PRESCRIBING MANUAL

OF THE

GHILDREN'S HOSPITAL

OF

PHILADELPHIA.

The doses of all the formulæ are proportioned for nildren who have reached the age of two years, one teasonful ([3]) being the dose of each preparation. The following table represents the doses for different es, apportioned according to this standard:

R

No. 1.

Liquor Cinchonidiæ Sulphatis.

S.

For No. 1000. Teaspoonful four times daily.

Date.

Signature.

Unless otherwise directed, the apothecary will always dispense three fluid ounces (f 3 iij) of the fluid preparations.

For convenience and certainty of measurement, when the dose to be prescribed is less than one fluid drachm (say f 3 ½ or f 3 ss), the apothecary will dilute a proportional quantity (say f 3 vj or f 3 jss) of the mixture with sufficient of some suitable menstruum to raise the amount to three fluid ounces (f 3 iij), and the patient will take of this one teaspoonful (f 3)).

MEDICINES

FOR

INTERNAL ADMINISTRATION.

LIQUORES.

Antiperiodic-

1.—Liquor Cinchonidiæ Sulphatis.

R	Cinchonidiæ sulphatis	gr. j
-	Acidi sulphurici diluti	m j fz j
	Aquæq, sad.	13)

M.—Dose, f 3 j, three times a day.

2.—Liquor Quiniæ Sulphatis.		
Ŗ	Quiniæ sulphatis	
N	ervous Sedativo—	
	3.—Liquor Potassii Bromidi.	
Ŗ	Potassii bromidi	
	M.—Dose, f 3 j, diluted, three times a day.	
	MISTURÆ.	
A	iterative—	
	4.—Mistura Arsenicalis.	
Ŗ	Infus. gentianæ comp	
	M.—Dose, f 3 j, three times daily.	
	5.—Mistura Forro-Arsonicalis.	
R	Liquor arsenici chloridi	

B. Syr. ferri iodidi	M v fg j
7.—Mistura Hydrargyri Chioridi	Corrosiv
R Hydrargyri chloridi corrosivi	gr. 1-96
Syrupi	f 3 ss
6.—Mistura Oloi Morrhu	180.
R Olei morrhuæ	fg ss
Syr. calcii lactophosphatis	m xv fg i
M.—Dose, f 3 j, three times daily.	-0,7
9.—Mistura Olei Morrhuæ (Comp.
R Olei morrhuse	fg ss
Syr. calcii lactophosphatis Syr. ferri iodidi	m∏ xv m∏ iv
Liquor calcisq. sad.	fgj
M.—Dose, f 3 j, three times a day.	
10.—Mistura Potassii iod	idi.
R Potassii iodidi	gr. j
Syr. sarsap. comp	fg ss

Antacid_	HOSPITALS.
R. Liquor calcis Aquæ cinnamo M.—Dose, f 3 j, e	Mistura Antacida.
R Magnati	a Magnesii Carbonatie
R Sodii bicarbonatis.	Sodii Bicarbonatia
M —Dose, [3], every	three hours.
Sodii bicarbonatis	Sodii et Gentianæ.

An	theimintic—		
	16.—Mistura Kameelæ Co	omp.	
Ŗ	KameelæSyr, acaciæ	gr. v f z ss	
1	Misce, et adde —	•	
ľ	Oleoresinæ filicis maris	m xv f ζ j one day.	
	17.—Mistura Olci Chonop	odii.	
1	Ol. chenopodii Pulv. acaciæ Syrupi Aquæ cinnamomi Q. sad. M.—Dose, f 3 j, three times a day for cowed by a laxative.	m iv gr. x f 3 ss f 3 j	>
	18.—Mistura Olei Chenopodi	i Comp.	
B.		m iv fg ss m 14 fg j	
	19. – Mistura Olooresinæ F	ilicis.	
B.	Oleoresinæ filicis	m xv m xx f z j	
foll	owed by a laxative if necessary.	iic day, to b	4

m. ii

mχ

Antiperiodic-20.-Mistura Ouinia Suiphatis. Ouiniæ sulphatis..... gr. i Acidi sulphurici dil m. i fg ss fg j Syr. zingiberis..... Aquæ.....q. s...ad. M.—Dose, f z j, three times a day. Antispasmodic-21.—Mistura Bolladonna et Aluminis. Ext. belladonnæ..... gr. I-24 Pulv. aluminis..... gr. ij Syr, zingiberis..... Syr. acaciæ..... m, xx M.—Dose, f z j, every three hours. Antipyretic-22.—Mistura Acidi Salicylici. R Acidi sallicylici..... Sodii bicarbonatis......āā... gr. j m xlviij Aquæ...... M. et adde-

Tr. aurantii cort.....

Glycerinæ

M.-Dose, fgj, p. r. n,

Astringont—

23.—Misture Acidi Sulphurici.

Ŗ.	Acidi sulphurici dil Liquor morphiæ sulph	m ij m viij
	Elix. Curagoæ	m, v
	Aquæq.sad.	fg j
N	M.—Dose, f z j, every three or four hou	rs.

24.—Mistura Argonti Nitratis.

R	Argenti nitratisSyr. acaciæ	gr. 1-64 M. x
	Aquæq. sad.	fgj
V	1.—Dose, f 3 j, every three hours.	

25.—Mistura Bismuthi.

Ŗ.	Bismuthi subnitratis	gr. v m. ss
	Syrupi	m, x
	Mist. cretæq. sad.	
ľ	1.—Dose, f z i, every three or four hou	rs.

26.—Mistura Cretæ Comp.

· B	Tr. opii camphorat	
	Tr. krameriæ	m v
	Sp. lavandulæ comp	m, ij
3	Misturæ cretæq. sad.	fʒj
	M.—Dose, f Z 1, every three or four hou	rs.

	27.—Mistura Crota et Rh	el.	
R	Syr. rhei aromat	m,	xv
	Syr. acaciæ	m.	
	Misturæ cretæq.sad.	fz	j
1	M.—Dose, fgj, every three or four hou	rs.	•
	28.—Mistura Rhei Com	n.	
R	Syr. rhei aromat		
Ļĸ	Sodii bicarb	gr.	XV
	Tr. opii camphorat	m,	
	Aq. menth. pipq. sad.	ſz	
,		•	,
ı.	M.—Dose, f 3 j, every three or four hou	rs.	
	29.—Mistura Zinci Oxid	li.	
R	Zinci oxidi	gr.	1/
	Glycerinæ	-	
	Syr. acaciæ āā	m	x
	Aquæq. sad.	f Z	j
1	M.—Dose, f 3 j, every four hours.		
Di	aphoretic—		
	30.—Mistura Antifebrill	B.	
R	Spt. ætheris nitrosi		
	Syr. limonis	m	
	Liquor ammonii acetatq. sad.	f Z	j
1	M.—Dos, f 3 j, every three or four hour	s.	
	31.—Wistura Potassii Citr	atis.	
R		m.	
134	Spt. setheris nitrosiq. sad.	f Z	
		U	j
į.	MDose, f z j, every two or three hour	·s,	

	32.—Mistura Potassii Citratis Comp.
R	Potassii citratis gr, ij s;
-	Spt. ætheris nitrosi m x
	Syrupi m xx
	Aquæ cinnamomiq.sad. f g j
	M.—Dose, f z j, every two or three hours.
Di	gestant—
	33.—Mistura Pepsinæ.
Ŗ	
	Acidi muriatici dil m iij
	Glycerinæ m v
	Aquæq.sad. f3 j
me	M.—Dose, f 3 j, three times a day in water, after eals.
DI	uretic—
	34.—Mistura Digitalis.
R	Potassii acetatis gr. ij ss
	Tincturæ digitalis m j ss
	Syr. scillæ m ij ss
	Syr. zingiberis m x
	Aquæq. sad. f 3 j
	M.—Dose, fgj, every two or three hours.
	35.—Misture Ferri Acetatis.
R	Tinct. ferri chloridi m, ij
~	Acidi acetici dil m. iv
	Liq. ammon. acetatis m. xl
	Syrupiq. sad. f 3 j
	M.—Dose, f 3 j, three or four times a day.

AMERICAN HOSPITALS.

	ERICAN HOOM	
36	stura Potassii Ac	LS.
R Pot	Stura Pot	
Cotassii acetati	- Classii Ac	ofor
Spt. juniperio	S	otatis.
Spt. ætherie	stura Potassil Ac is	gr. ijs
SVIII	UOS1	m - 1) S
* FOIDS	******	m v
M Do	***************************************	m x
Dose, f3j, e	very three or s	m xv
	three or four h-	13 j
Expector	very three or four hour	S.
SrMist. A-		
R	onii Carbonatis et	
R Ammonii carbonal Syr. tolutani Syr. senega	onatis et	
Syr. tolutani	US	senega
		gr. j
Syr. tolutani	***************************************	m x
M. Dose f	n	liij
3), every	two '9. Sad. fo	3 j
M.—Dose, f3j, every	or three hours	, ,
30 -		
Jo. Mistura	Ammonii Chloridi	
Ammonii ali	Ammonii Chi	
Ext. glycomidi	Unioridi	
Ammonii chloridi Ext. glycyrrhizæ fld Syrupi		**
	ur.	

	40Mistura Ipecacuanha	Comp.
R.	Potassii citratis	gr. iij m iij m ss. m xv f z i
	41.—Mistura ipecacuanhe et	Scilia.
B.	Syr. ipecac	m v m iv fg j
	42.—Mistura Pruni Virgini	anæ.
R N	Acidi sulphurici dil	m j m ss f 3 ss f 3 j
	 43.—Mistura Scillæ Con	ân.
R		.h.
-	Syr. scillæ	m ijss m iv fgj irs.

La	xative and Cathartic—		
	44.—Mistura Aloes et Fe	rri.	
I&	Tr. aloes et myrrhæ Ferri sulphatis exsiccat Syr. rhei aromat	m gr. f z	1.6
	45.—Mistura Magnesii et	Opii.	
13.	Magnesii sulphatis	gr. m m f 3	v ss x
	46.—Mistura Olei Ricin	i.	
R	Olei ricini	f g m	SS 1/8
	47.—Mistura Podophyl	1	

H. Inf. sennæ	m xv fg j
Nervous Sedative	
B. Tr. belladonnæ	m ijss m xv fgj
B1.—Misture Chiorel. Hyd R Chlorel. hydrat	gr. ijss
B. Chloral. hydrat	Bromidi. gr. ij gr. v f 3 j
Stimulant— 53.—Mistura Ammonii Carbo R Ammonii carbonatis	gr. j m. xx f 3

	54.—Mistura Spiriti Vini Q	ailici.
Ŗ	Spiriti vini gallici	m xv
	Syrupi	mχ
	Aquæq. sad.	fg j
	M.—Dose, fgj, every three or four hou	rs.
T	onic—	
	55.—Mistura Cinchonidiæ (Comp.
R	Cinchonidiæ sulphatis	gr. j
·	Acidi muriatici dil	Μj
	Aquæ	m xxx
	Ft. sol. et adde	
	Potassii chloratis	gr. ij ss
	Tinct. ferri chloridi	Μij
	Syr. zingiberisq. sad.	ſgj
	M.—Dose, fgj, four times a day or ofte	ner,
	_	
	56Mistura Ferri et Ammon.	Citratis.
R	Ferri et. ammon. citratis	gr. j
	Tr. gentian. comp	ηx
	Spt. lavand. comp	щν
	Syr. limonisq. sad.	fžj
	M Dose, f 3 j, three times a day.	
	57.—Mistura Ferri Cen	p.
R	Tinct. ferri chloridi	
	Acidi acetici	m ijss
	Liquor ammonii acetatis	m xl
	Potassii chloratis	gr. ij ss
	Syr. zingiberisq. sad.	fдj
	M.—Dose, f 3 j, every three or four hou	ırs.

	58.—Mistura Ferri et Potassii	Tartratis.
Ŗ	Ferri et potassii tart Tinct. gentianæ comp Spt. lavandulæ comp Syr, limonisq. sad. M.—Dose, f z j, three times a day.	gr. j ss m xv m v f z j
	59.—Mistura Ferri et Qui	niæ.
Ŗ		gr. ss m ij m xx f g j
	80 — Mistura Nucis Vemi	l
R		m j m v fgj
	81.—Mistura Potassii Chie	ratis.
Ŗ	Potassii chlorat	gr. ij m ij f 3 ss f 3 j y, in water.

PULVERES.

Antacid—
62,—Pulvis Bismuthi et Sodii.
R Bismuthi subcarbonatis
Sodii bicarbonat
M. et ft. chart. No. j.—Dose, one powder thrice daily.
63.—Pulv. Ipecac. et Cretæ Præperat.
R. Pulv. ipecac. comp gr 1/3
Cretæ præparatæ gr. iij
M. et ft. chart No. j.—Dose, one powder every six hours.
nours.
Antheimintie-
64.—Pulvis Santonini.
R Santonini gr. ss
Sacchari gr. j ss
M. et st. chart. No. j.—Dose, one powder morning and
evening, the second to be followed by a purge.
Astringent—
65.—Pulvis Bismuthi.
R Bismuthi subnitratis gr. j
Sacchari gr. ij
M. et ft. chart. No. j.—Dose, one powder.
66.—Puivis Bismuthi Comp.
R Bismuthi subnitratis
Pulv. aromatici
M. et ft. chart. No. j.—Dose, one powder every four
or six hours

	67.—Pulvis Bismuthi et Acid. Tannici.
Ŗ	Bismuthi subnitratis gr. j
	Acid. tannici gr. ss
	I. et ft. chart. No. j.—Dose, one powder every six
hou	rs
	00 0 1 1 0 01 0 11 14 0 11
_	68.—Pulvis Bismuthi et Opii.
ĸ	Bismuthi subnitratis gr. iij Pulv. opii gr. 1-10
M	I. et ft. chart. No. j.—Dose, one powder every six
hou	
69	Pulvis Bismuthi, Ipeeac. ot Ac. Tannici
Ŗ	Pulv. bismuthi et ac. tannici (No. 67) gr. j ss Pulv. ipecac. comp gr. ½
N	I. et ft. chart. No. j.—Dose, one powder every six
hou	rs.
	70.—Pulvis Plumbi et Opii.
Ŗ.	Plumbi acetat
	Sacchari gr. ij
N hou	f. et ft. chart. No. j.—Dose, one powder every six rs.
	71.—Pulvis Tostæ Præparat. et Opii.
Ŗ	Testæ præparatæ gr. iij Pulv. ipecac. comp gr. ½
N	M. et ft. chart. No. jDose, one powder every six
hou	

Digestant-	_	
72	Pulvis Bismuthi et Pep	sinæ.
R Bismuth	i subnitratissaccharat	
M. et ft. cl	hart. No. j.—Dose, one powd	er three times
73	.—Pulvis Pepsinæ et Sc	dit.
R Pepsinæ Sodii bio	saccharatcarbonatis	gr. ij gr. j
M. et ft. ch daily.	nart. No. j. —Dose, one powd	er three times
	and Cathartic—	
	Pulvis Hydrarg. Chlorid	d. Mit.
Sacchari		gr. j gr. ij
M, et ft. ch	art. No. j.—Dose, one powde	e r.
75	Puivis Ipecacuanha et	Rhei.
R Pulv. ip	ecacuanhæ	gr. ¼
Pulv. rhe	eieirbonatis	gr. j
	art. No. jDose, one powde	· .
111 01 111 01	——————————————————————————————————————	i timiec dany.
76.	.—Puivis Rho i ot Gentia	nas.
	pulv	gr. j gr. ij
	art. No. jDose, one powde	
a day.	• •	•

77.—Pulvis Rhei et Magnesii.

R.	Pulv. rhei	gr. i
	Magnesii carbonatis	gr. v
	Pulv. zingiberis	gr. 1-6
1	M. et ft. chart. No. j Dose, one powder	thrice daily.

ENEMATA.

78.—Enema Argenti Nitratis.

R.	Argenti nitratis	gr. ¼	
	Bismuthi subnitratis	gr. xx	
	Tinct. opii deodorat	m iij	
	Mucil. acaciæ	ſg j	
	Aquæq. sad.	ſžij	
3	1.—Dose, fzij, injected into bowel	twice a	day.
	ike well.		•

79.—Enema Opii.

R	Tinct. opii	m, iij
	Mucil. acaciæ	fg i
	Aquæq. sad.	f z j fzij
N	1.—Dose, f z ij, injected into bowel tw	

PREPARATIONS

FOR

EXTERNAL USE.

The various fluid preparations embraced in this list are to be kept and dispensed in *colored* and *roughened* bottles, labeled "For External Use."

GARGARISMATA.

B.	Acidi tannici	nnici. 3 j f3 j f3 j
R.	81.—Gargarisma Alumir Pulv. aluminis	gr. xv m x f z j f z j
B.	82.—Gargarisma Potassii Cl Potassii chloratis	gr. xx fz j fž j

	i.—Gargarisma Potass. Chior Carbellol.	at. et Ack	
	Acidi carbolici	gr. ij f ž j	
84	.—Gargarisma Potass. Chlore	nt. et Ferr	ı.
B.	Tinct. ferri chloridi	m x f 3 j	
	85.—Gargarisma Scdii Bo	ratis.	
R •1	Sodii boratis	gr. xv f 3 ij f 3 j	
	GLYCERITA.		
	86.—Glyceritum Iodini	ii.	
R.	Iodinii	gr. v [3 j	
	87.—Glyceritum Tr. Forri Cl	hiorisi.	
R.	Tr. ferri chloridiq. sad, Glycerinæq. sad, M.	f 3 j	

LINIMENTA.

R	88.—Linimontum Aconiti C	emp.
	Chloroformiāā Liniment, saponisq. sad.	m xv fz j
1	M.	
	89.—Linimontum Ol. Suc	ei ni.
R	Ol. suecini	
	Ol. olivæāā	f 🖁 ss
1	M.	
_	90.—Linimentum Sapen. V	
Ŗ	Saponis viridisAlcoholis	g j f ž i
7	M.	ر و.
-		
	LIQUORES.	
	S1.—Liquor Acidi Boraci	ci.
R	Acidi boracici	gr. xv
	Aquæ destillatæ	gr. xv f 🖁 j
1	M.	
	92.—Liquor Acidi Tanni	ici.
R	Acidi tannici	gr. viij
-	Aquæ destillatæ	gr. viij f Z j
1	M.	

	93.—Liquor Aluminis.		
Ŗ.	Aluminis	gr. f 🕇	iv
	Aquæ rosæ	f 3	j
1	M.		
	94.—Liquor Argenti Nitre	lis.	
Ŗ.	Argenti nitratis	gr.	SS
_	Aquæ destillatæ	f 3	J
2	M		
_	95.—Liquer Atropiæ.		
ĸ	Atropiæ sulphatis Aquæ destillatæ	gr. f 3	1V
,	M.	13	J
1	v.,		
	96.—Liquor Hydrargyr	.	
D.			1.6
1,1	Aquæ destillatæ	gr. f 3	i
1	м. •	J	•
	97.—Liquor Plumbi Subace	teti	s .
R	Liquor plumbi subacetatis	m	χV
	Aquæ destillatæq. sad.	ſξ	j
1	M.	Ŭ	
	98.—Liquor Sodii Borati	is.	
Ŗ		gr	x
	Aquæ destillatæ	fZ	j
]	М.		

LOTIONES.

	~~~~		
	99.—Lotio Acidi Sulphur	osi.	
Ŗ.	Acidi sulphurosi	fʒ ijss f₹ 1	
	Aquæq. sad.	fåj	
Y		•	
	100.— Letio Coccuil Indi	ici.	
R	Tinct. cocculi indici		
	Alcoholisāā	f 🕱 ss	
3.4		. 5 33	
IV.	I.—Sig. To be diluted.		
	101.—Lotie Sodii Hyposuip	hitis.	
R.	Sodii hyposulphitis	z ii	
	Aquæ destillatæ	3 ij 13 i	
M	<u>.</u>	0.	
	<del></del> 0- <del></del>		
	UNGUENTA.		
102	 2.—Unguentum Hydrarg. Amr	nonisti Di	11
ľ	Ung. hydrarg. ammoniat	3 ss	
	•	3 33	
M	l.		
	-		
10	D3.—Unguentum Hydrargyri I	litrat. Dii.	
	Ung. hydrarg. nitrat		
	Adipis	3 j 3 vij	
M	•	0 3	
-			

	104.—Ung. Hydrarg. Oxidi Flav. Dil.
R	Hydrarg. oxidi flav gr. viij
	Hydrarg. oxidi flavgr. viij Ung. petrolei
V	ſ.
10	5.—Unguentum Hydrarg. Oxidi Rub. Dii.
Ŗ	Ung. hydrarg. oxid. rub
	Ung. hydrarg. oxid. rub
N	1.
100	3.—Unguentum Hydrargyri et Belladonnæ
Ŗ	Ext. belladonnæ
	Ung. hydrargyri
_	
N	ſ.
	107.—Unguentum Picis Liquidæ Dij.
Ŗ.	
_	Adipis 3 ss
V	ſ.
10	8.—Unguontum Piumbi Subacetatis Dil.
Ŗ.	Cerat. plumbi subacetat
	Cerat. plumbi subacetat
	Aupts
V	<b>1.</b>
104	9.—Unguentum Sulphuris et Potass. Carb.
R	
ľķ.	
	Ung. sulphuris
1	Adipis 3 vj
•	12.

#### APPENDIX:

Containing those preparations of the U.S. Pharmacopaia most commonly employed in Children's Diseases, together with their doses for Children two years old. Aceta-Acetum opii..... m, ss-i scillæ..... m j-v Acida ---Acidum aceticum dil..... m xv-v carbolicum ..... gr. 1/8 " gallicum ..... gr. ss-ii hydrocyanicum dil..... 66 m 1/8-1/4 m 1-v " muriaticum dil..... " nitricum dil..... m j-iij nitromuriaticum dil..... m j-v " phosphoricum dil..... m. i-v " salicylicum ..... gr. j 66 sulphuricum dil..... m j-v " aromat..... m j-v " tannicum..... gr. ss-ij Ætherea... Æther ..... m ij-x Chloroformum..... m, j−v Ammonia.... Ammonii bromidum..... gr. j-v " carbonas ..... gr. ss-j

chloridum.....

gr. j-v

"

## OFFICIAL FORMULÆ

Acid. nitric. dil " muriatic. dil M.	
	<del></del>
Antimonium—	·
Antimonii et potassii ta	rtras gr. 1-48-14
Aquæ	
•	1)
Aquæ ammoniæ (Exter	nai)
" cinnamomi	f7 i-ii
" menthæ piperitæ	
" rosæ (External).	
-	
Argentum—	<del></del>
·	
Argenti nitras	gr. 1-64–1-32 ternal)
Bismutham—	
Bismuthi subcarbonas	gr. j <b>–v</b>
" subnitras	gr. j <b>–v</b> gr. j–v
Calcium—	
Calcii lactiphosphatum.	gr. j–ij
Creta præparata	
Testa '··	gr. ij–x
Corata (External)—	<del></del>
Ceratum canth.	Ceratum resinæ comp.
" cetacei.	" sabinæ.
" ext. canth.	" saponis.
" plumbi sub.	" zinci carbon.
" resinæ.	

## Charts (External)-

Charta cantharidis. "sinapis.

## Collodium (External)-

Collodium cum cantharide. Collodium flexile.

#### Confectiones-

Confectio sennæ..... gr. x-xx

#### Dececta-

Decoctum dulcamaræ		f <b>z</b> j-ij
"	hæmatoxyli	f z j-ij
44	hordei	fg j–ij fg j–ij fg j–iv
"	quercus albæ (External)	0,
66	uva: ursi	fz i-i:

## Emplastra (External)—

Emplastrum assafœtidæ.

- " belladonnæ.
- " hydrargyri.
- " opii.
- " picis burgundicæ.
- " cum cantharide.
  " resinæ.
- " saponis.

Extracta-		
Extractur	n belladonnæ	gr. 1-32-1-16
66	cinchonæ	gr. j–iv
44	colocynthidis c	gr. <b>¼</b> -j
"	gentianæ	gı. ss–j
. 66	glycyrrhizæ	gr. j-v
",	hæmatoxyli	gr. j-iv
66	hyoscyami	gr. 1-16-1/
44	krameriæ	gr. ss-ij
"	nucis vomicæ	gr. <b>1-48–1-2</b> 4
**	taraxaci	g <b>r</b> . ij–x
Extracta F	luida—	
	a buchu fluid	m ij_v
######################################	cimicifugæ fluid	m iv-viij
66	ergotæ fluid	m j-ij
60	jaborandi fluid	m, x
**	pruni virg. fluid	. m. x
"	rhei fluid	m j_v
66	sennæ fluid	m z-xxx
"	spigeliæ et sen. fluid	fzi
"	uvæ ursi fluid	m lj_v
66	valerianæ fluid	mŭij–x
		• •
Forrum—		
	<b>18</b>	gr. ss–ij
	mmonii citras	gr. <b>ss</b> –ij
	otassii tartras	gr. ss–ij
	uiniæ citras	gr. ss–ij
	as	gr. ss-ij
	ophosph <b>as</b>	gr. 🔏 –j
	carbonas	gr. j–ij
	has exsiccata	gr. <b>¼</b> −j
Ferrum re	dactum	gr. s <b>s</b> –ij

#### Glycerita (External)-Glyceritum acidi carbolici. gallici. " tannici. 46 picis liquidæ. sodii boratis Hydrargyrum-Hydrargyri chloridum corros...... gr. 1-100-1-32 mite..... gr. 1-24-ss Hydrargyrum cum creta..... gr. ss-jss Infusa --Infusum buchu..... f **g** j-ij f **g** j-ij calumbæ..... " digitalis..... m x-fg ss " gentianæ comp..... f g ss-j " jaborandi..... f z j-ij f z j-ij f z ss-i " juniperi..... " lini comp..... ss-ii f **g** ss-j f **g** j-ij " sennæ..... " spigeliæ..... Linimenta (External)-Linimentum aconiti. ammoniæ. " calcis. " camphoræ. " cantharidis. " chloroformi.

plumbi subacetatis.

saponis

terebinthium

66

"

"

Liqueres		
Liquor  " " " " " " " " " " "	ammonii acetatis	f3 ss-j m, j-iij m, ss-j f3 j-f3 iij m, ss-j f3 ij-f3 ss m, j-iij f3 ss-j
magnesi	<b>um</b> -	
Magne Magne	sii carbonas	gr. v-9 ij gr. v-9 ij
		:
<b></b>	•••••••••••••••••••••••••••••••••••••••	gr. xx-3 j
Mellita (i	External)—	
Mel de	espumatum.	
	sæ.	
" so	dii boratis.	
Mistura-	_	
	a ammoniaci	£= ::
Mistur	ammoniaciamygdalæ	fg ss-ij
**	assafœtidæ	13 1-13 1
44	chloroformi	(2 i-ii
44	cretæ	f3 j-f3 j f3 j-ij f3 j-ij f3 ss-j
**	ferri comp	f <b>z</b> j–ij
"	glycyrrhizæ comp	m xv-fz ss
"	potassii citratis	f z ss-j

Morphia—	
Morphiæ acetas	gr. 1-48-1-30 gr. 1-48-1-30 gr. 1-48-1-30
Mucilagines—	
Mucilago acaciæ	ad lib. ad lib. ad lib. ad lib.
Olea— Oleum chenopodii	m ij-iv m ss-j m xv-f z j f z ji-j f z ss-ij m ij-v
Oleoresinæ	
Oleoresina filicis	m v-xx
Opium	gr. 1-10-1 8
Pepo	<b>3</b> j-ij
Phosphorus—	gr. 1-160 1-100
Plumbum— Plumbi acetas	gr. 1.8–1-6

Potassium –	
Potassii acetas	gr. ij-v
" bicarbonas	gr. ij–v
" bitartras	gr. x-xv
" bromidum	gr. ij–v
" chloras	gr. ij–v
" citras	gr. ij-v
" et sodii tartras	gr. xv-3 j
" iodidum	gr. ss-iij
" nitras	gr. jss-viij
Pulveros—	
Pulvis aromaticus	gr. j–ij
" ipecacuanhæ comp	gr. ¼-jss
•	
Quinia —	
Quiniæ sulphas	gr. ss-iv
" valerianas	gr. ¼–ss
	· ,.
Resina-	
Resina jalapæ	gr. ¼ −ss
" podophylli	gr. 1-64-1.45
" scammonii	gr. 🏏 – jss
Rheum	gr. ss–ij
Pottlere (Versele)	
Rettiora (Kameola)—	gr. v–xv
Santoninum	gr. ¼–ss
	5 /4
Scammonium	gr. j–ij
Sensa—	gr. iij-v
	81. m-1

Sinapis (Emetic)—	gr. viij–xx
Sodlum—	
Sodii acetas	gr. ij–v r. 1-160–1-64 gr. ij–v
*****	
Spiritus—	
Spiritus ætheris comp	m ij-x m v-xx m ij-v m ss-iv m j-ij m v-f z j m v-xx m v-xx m ss-ij m v-f z j
Strychnia—	
Strychniæ sulphas, gr.	1-150-1-100
Bulphur	
Sulphur præcipitatum	gr. v-xv

Syrupi—			
Syrupus	acaciæ	ad	lib.
<i>"</i> " •	ferri iodidi	m.	ij-v
66	ipecacuanhæ		íj–fgj
66	krameriæ	m.	xx-iz
66	lactucarii	m	xx-13 j
66	limonis	ad	lib.
**	pruni virginianæ	m,	xv-f3ss
"	rhei	fg	
"	" aromat	fž	ss-j
**	sarsaparillæ comp	Ψ	
66	scillæ	m	ij-vj
••	" comp	m	ij–vj
66	tolutani	m	ij_vj
"	zingiberis	m	V-X
Tinctur <b>æ</b> -	-		
Tinctura	belladonnæ	m	j–ij
"	calumbæ	m	iij–xv
"	cannabis	m	j <b>–</b> ij
"	cardamomi comp	m	v-xv
"	catechu	m.	v-xv
46	cinchonæ comp	m	X-XV
44	cinnamomi	m	ij–xv
"	colchici	m	j–iij
44	Digitalis	m	ss-iij
66	ferri chloridi	m	j-v
"	gentianæ comp		v-xv
44	guaiaci ammon	m	VXV
66	hyoscyami	m	j–iv
46	iodinii comp	m	j-iv
46	kino	ηŭ	ij–xx
"	krameriæ	mį	ij-xx
"	nucis vomicæ	m.	ss-ij

Tinctur	a opii	m j–iij m v–xx m j–iij m j–v m v–xv m ij–vij
Unguenta	(External)—	
	tum acidi carbolici.	
0 8 0	" tannici.	
"	aquæ rosæ.	
46	belladonnæ.	
66	benzoini.	
68	cantharidis.	
**	creasoti.	
**	gallæ.	
66	hydrargyri.	
**	" ammoniati.	
4	" iodidi rubri.	
66	" nitratis.	
"	" oxidi flaxi.	
"	" " rubri.	
46	iodinii.	
46	" compositum.	
46	mezerei.	
"	picis liquidæ.	
"	plumbi carbonatis.	
16	" iodidi.	
**	potassii iodidi.	
"	stramonii.	
"	sulphuris.	
"	" iodidi.	
"	veratriæ.	
16	zinci oxidi.	

#### Vine-

/inum	ergotæ	m v-x
"	ipecacuanhse	gtt. ij-viij
"	opii	m ss-j
"	rĥei	m v-fz ss
	xericum	m x—fz i
	portense	m. v-fz i

## Zincum-

Zinci	oxidum	gt.	1/4-ss
"	sulphas	gr.	1⁄4-v
"	valerianas	er.	1/4-ss

## Zingiber---

Pulv.	zingiberis	gr.	1-6-i	ij

## NON-OFFICINAL REMEDIES.

Acidum salicylicum	gr. ss-iss
Pepsina saccharata	gr. j-v
Pulv. glycyrrhizæ comp	gr. j <b>–v</b> gr. iv–viij
Sodii bromidi	gr. j-v
Sodii salicylici	gr. ss-j ss

#### POISONS AND THEIR ANTIDOTES.

Acids, Mineral.—Alkalies in dilute solution; lime; whitewash; magnesia; soap.

Quickness of administration essential.

Avoid emetics and stomach-pump.

Acids, Vegetable—Soap; soda or potassa in dilute solution; except for oxalic acid, for which give lime, whitewash, or chalk.

Aconite.—Emetics; stomach-pump; tannic acid; digitalis; laudanum; warmth; stimulation; recumbent

position.

Alkalies, and their salts. - Vinegar; weak acids; ou, freely; opium.

Alum .- Albumen.

Antimony, and its salts.—Tannic acid; opium; external warmth.

Arsenic, and its salts.—Emetics; stomach-pump; recently made hydrated sesquioxide of iron; magnesia; dialysed iron; oil or fat.

Atropia. See Belladonna,

Belladonna.—Emetics; stomach-pump; tannic acid; morphia hypodermically.

Bismuth, and its compounds .- Albumen; milk.

Bromine. - Soap; oils.

Cannabis Indica.—Emetics; lemon juice; quiet; if exhaustion, stimulants.

Carbolic Acid.—Stomach-pump; solution of saccharate of lime; hot and cold douche.

Chloral and Chloroform.—Alternate hot and cold douche; artificial respiration; cardiac stimulants.

Cocculus Indicus.—Emetics; stomach-pump; at first, give opium; chloral; chloroform; later, stimulants.

No chemical antidote.

Colchicum.—Emetics; stomach-pump; tannic acid; opium; stimulants.

Conium.—Emetics; stomach-pump; artificial respiration; tannic acid; opium.

Copper, and its salts.—Albumen; milk; calcined magnesia; yellow prussiate of potash.

Croton Oil, and other drastic purgatives.—Emetics; albuminous drinks; bismuth; external heat.

Cyanide of potassium .- See Hydrocyanic acid.

Digitalis.—Emetics; stomach-pump; tannic acid; stimulants.

Gelsemium .- Same as Aconite.

Hydrocyanic acid.—Alternate hot and cold douche; intravenous injection of aqua ammoniæ; atropia hypodermically.

Hyoscyamus.-Same as Belladonna.

Iodine.—Starch, freely; if this cannot be obtained, then soap.

Iron, Salts of-Soap; dilute alkalies; albumen.

Lead, Salts of—Emetics; stomach-pump; alkaline sulphates; soap; albumen; milk; purge.

Mercury, Salts of-Emetics; albumen; milk.

Opium.—Stomach-pump; atropia hypodermically; coffee; flagellation; artificial respiration; electricity.

Phosphorus.—Emetics; sulphate of copper in small doses; crude oil of turpentine; stomach-pump; avoid oils or fats.

Potash Salts .- No distinct antidote.

Santonin.—Emetics; purges; stimulants; artificial respiration.

Silver, Salts of-Common salt, freely; albumen.

Soda Salts .- No distinct antidote.

Stramonium.-Same as Belladonna.

Strychnia.—Emetics; stomach-pump (at first only); tannic acid; chloral; bromide of potash; ether; chloroform; rest.

Veratrum viride. - Same as Aconite.

Zinc, Salts of—Albumen; carbonate of soda; milk, freely,

#### INFANT DIET.

Diet from Birth to Age of Six Months.

(Five to six meals in 24 hours.)

MILE, CREAM, LIME-WATER, ARROWROOT WATER.

Take four tablespoonfuls of each, place in bottle with a lump of sugar, warm the whole by standing bottle in warm water.

Arrowroot Water is made by taking one and a half teaspoonfuls of arrowroot, rubbing it down until smooth, with a tablespoonful of cold water, and then adding one pint of boiling water, stirring all the time.

Diet from Six to Twelve Months Oid.

(Five meals in the day.)

FIRST MEAL, 7 A. M.

Mellin's food, one tablespoonful, or Flonr Ball grated, one or two teaspoonfuls.

Hot water, four tablespoonfuls.

Warm milk, enough to make half a pint.

Dissolve the Mellin's food or rub down the grated Flour Ball in the hot water by stirring, then add the milk, and mix thoroughly.

SECOND MEAL, 10.30 A. M. THIRD MEAL. 2 P. M. A breakfast cupful of milk with four tablespoonfuls of lime-water,
FOURTH MEAL, 5.30 P. M. Same as first.

FIFTH MEAL, 10.30 P. M. Same as irst. Fifth MEAL, 10.30 P. M. Same as second.

Diet for a Child One Year Old.

(Five meals in the day.)

FIRST MEAL, 7 A. M.

Two teaspoonfuls of grated Flour Ball in half a pint of milk.

SECOND MEAL, 10.30 A. M.

Half a pint of milk with four tablespoonfuls of limewater.

THIRD MEAL, 2 P. M.

The yolk of one egg beaten up in a teacupful of milk.

FOURTH MEAL, 5.30 P. M. Same as the first.

FIFTH MEAL. Same as the second. FOR DRINK—Filtered water.

ni e dilla ii

Diet for a Child Two Years Old.

Breakfast, 7.30 A. M.

Milk.

The lightly boiled yolk of an egg. Thin bread and butter (bread one day old).

LUNCH, II A. M.

Milk.

A thin slice of bread and butter.

DINNER, 1.30 P. M.

Beef-tea, or small piece of minced roast beef or mutton. One well-mashed potato, moistened with gravy.

Rice and milk.

#### OF AMERICAN HOSE

SUPPER, 6 P. M.

Milk.

Bread and butter.

FOR DRINK-Filtered water.

Diet in Mucous Disease for a C

BREAKFAST, 7.30 A. M.

Milk, with lime water; four water to each tumblerful of mill The lightly-boiled yolk of well-toasted bread, or stale brea DINNER, 12 M.

A mutton chop without fat roast beef or mutton, occasional Stale bread or toast.

SUPPER, 7 P. M.

Milk and lime-water. Stale bread or toast. FOR DRINK-Filtered water.

Starch food to be avoided a

#### BARLEY WA

-0-

To prepare barley water, put t washed pearl barley with a pint pan, and boil slowly down to tw

BEEF-TE

Take one pound of lean beet with its juice, into an earthen ve tepid water, and let the whole st well, squeezing all the juice from fire, and slowly raise just to the bo briskly all the time. Season with sa

In administering this, always be

sediment.

#### FLOUR BALL.

Take one pound of good flour—unbolted, if possible—tie it up very tightly in a pudding-bag; put it in a pot of boiling water early in the morning, and let it boil till bed time, then take it out and let it dry. In the morning peel off from the surface, and throw away, the thin rind of dough, and with a grater, grate down the hard dry mass into a powder. To use this, take from one to two teaspoonfuls of the powder, rub it down till smooth with a tablespoonful of cold milk, and add a tumblerful of hot milk, stirring it well all the time.

#### LIME-WATER.

Take a piece of unslacked lime, as large as a walnut, put it into two quarts of clear water, stir about thoroughly, allow to settle, and use only from the top.

#### WINE WHEY.

Boil in a pint of fresh milk; while boiling, pour in a small tumblerful of sherry wine (eight tablespoofuls), bring it to the boil a second time, being careful not to stir it; as soon as it boils set it aside until the curd settles, and pour off the clear whey.

# PENNSYLVANIA HOSPITAL

۸r

# PHILADELPHIA.

# St. John Long's Liniment.

1328	1 29 11 4
	ノること

#### Lead-Water and Laudanum.

Face Maret and Faceur	
Take of—	
Laudanum	oz. 4
Ganlard's extract	oz. 4
Water, to make one-half gallon.	-
Mix.	
The state of the s	
Solutions of Carbolic Ac	id.
2 per cent., 21/2 per cent., 4 per cent.	and 5 per cent.
(To make two per cent. solution, add a crystals and heat to make it liquid, and pu acid (by measure) to fifty ounces of water crystals liquid, set bottle in hot water; the water to keep it liquid.)	t one ounce of er. To render
Solution of Boracie Acle	d.
R Acid. boracici	gr. xv
Aquæ puræ	gr. xv
M.—Used mostly for sore eyes.	200
400 000 000	
Carbolized Oil.	
Take of—	2000
Calvert's carbolic acid, No. 4	oz. I
Olive oil	oz, 15
Mix.	
Aperient Pills.	
R Ex. colocynth comp	gr. xxiv.
Ex. nucis vom	gr. iij
Ex. belladonnæ	gr. j
M Ft. pil. No. xij Sig. One or two	
The Property of the or the	No. of Contract of

ΛÞ	AMPRICANT	HOSPITALS.
Or	AMEKICAN	HOSPITALS.

105

# Niemeyer's Pills.

R	Quinise sulph		gr. xi	j
	Pulv. digitalis	• • • • • • • • • • • • • • • • • • • •	gr. vj	
	Pulv. opii	•••••	gr. iij	
3	M.—Ft. pil. No. xij.—Sig.	One, three	times a	day.

#### Diarrhosa Mixturo.

re oi—			
Chalk mixture	oz.	4	
Paregoric	oz.	2	
Tr. catechu	oz.	2	
Mix -Dose two teaspoonfuls			

#### Chierate of Patassa Solution.

$\mathbf{R}$	Potass.	chlor	gr.	xxx.
		***************************************	3	j
T	Dissolve	For throat trouble		

# **Compound Gontlan Mixture.**

Ŗ.	Ferri cit. et ammon	<b>3</b> ij
	Fl. ex. gentian	ž vi
	Spt. lavand. comp	Ŏi
	Alcoholis	O ss
	Syrup. simp	Z xxiv
	Aquæad	Ċj
1	M.—Use as a tonic.	-

#### Basham's Mixture.

R	Sal. ammon. acet	3 xxx
	Acid. acetas	3 xv
	Tr. ferri chlor	Z xxv
	Alcoholis	<b>3</b> x
	Syr. simplicis	₹ xx
	Aquæ	3 xv

M.—Dose, tablespoonful three times a day.

(Note.—It will be noticed that the proportion of alcohol in the above is greater than in the officinal formula. This makes it keep better in a warm place.)

#### Solution Iodidi Potass.

ĸ	Potass iodidi	Э	j
	Aquæ cinnam		SS
٠,	Doce tablespoonful	C	a aal

M.—Dose, tablespoonful. Sometimes we use solution double strength.

# Solution Quinæ.

Ŗ	Sul. quiniæ	ξij ζjgr. iv m lxxx
	Acid sulph. c. p	
	Aquæ	one gallon

Put quinine in mortar, add half pint water, add acid and dissolve, then add remainder of water. This gives a solution of one grain to the ounce.

# JEFFERSON COLLEGE MOSPITAL

OF

#### PHILADELPHIA.

#### .lodotorm Cotton.

Take of-

Cotton gr. 70 Glyecrineminims 10	
Alcohol, (sp. gr. 796)	ct
Mix.—Dissolve the iodoform in the ether, add the gi cerine, mixed with the alcohol; saturate the cotton wi the above mixture, and dry by exposure.	
Koumys.	
Take of—	
Sugar dr. 6 Compressed yeast, (very fresh, size of pea).	
Mix.—Fresh milk, to fill champagne bottle. Co tightly; keep in temperature of about 80°, first twent four hours, and shake occasionally; then keep in co place (on ice) two days.	ty

106	AL FORMULÆ	
	owder—(Sajou's	B).
	.e.	gr. ss gr. xxx 3 iss 3 j
	él's Pill of Hydrarg. C	omp.
	Mass. drarg	gr. j gr. ss gr. ij gtt. ss
	S. W. Cross' Fover Mixtur	⊕.
	Liq. potassii citratis	3 ss 3 j gtt. ij gr. 1-16
M	—Sig. To be taken every three, four o	r five hour

# 

M. ft. sol. Ten minims contain four grains of ergitine.

# S. D. Gross' Antimony and Saline Mixture.

Antimonii et potass. tart	gr. ij ss
Magnesii sulphas	3 ij
Morphiæ sulphas	gr. i1/3
Aquæ dest	3 x
Syr. zingiberis	Žij
Tr. virat viridi or aconiti	ž jss

M.—Sig. Tablespoonful every two, three, four or six hours. Used in acute stages of gonorrhea and other inflammations.

#### Sojou's Douche.

(For Nasal Catarrh.)

Sodii salicylici	3 x
Sodii biboras	ž xv
Glycerini	🕇 ij ss
Aquæ	₹ xxx
r *C: 17 J1	

# M.—Sig. Use as a douche or spray morning and night

# Parvin's Carcinoma Solution.

R	Iodinii	<b>3</b> j
	Brominii	3 1
	Acid. carbolici	3 iv
	Alcoholis	žј

M.—Apply to parts, and follow with saturated solution of bicarbonate of soda, to neutralize any excess. To be used to destroy diseased tissue not removed by operation.

# __ Unguentum Bero-Glyceride.

Boro-glyceride (50 per cent)	3 j
Ung. petrolei	3 v
Mix with gentle heat.	

Used in conjunctivitis, and ulcerated cornea. May be used in ulcerated conditions in any part of the body.

#### Boro-Glyceride.

#### Take of-

Glycerine (50 per cent. solution) Boracic acid	gr.	92
Boracic acid	gr.	62

Stir with gentle heat until mixture crystalizes upon cooling; then add, while warm, 50 per cent. of glycerine by weight. This can be diluted with water to any desired strength. Used eztensively here, in 25 per cent. solution, for catarrh of nasal and pharyngeal membranes.

#### Mist. Guaiaci Comp.

#### (For Throat Troubles.)

R	Tr. guaiaci ammon	
	Tr. cinchon. compāā	3 ij
	Potass. chloras	ЗÍ
	Mel. despunat	3 vj
	Aquæq. s	3 vj 3 iij
	M.—Sig. To be used as a gargle and	swallowed.

## Lupuiin Mixture.

#### (Diuretic.)

12.	Lupulinæ	~ ii
1X	Uvæ ursi	2 :
	Sodæ bi-carb	3 1) 2 55
	Tr. opii camph	2 iv
	Aquæq. s	Όij

Make decoction of lupulin and uva ursi and add other ingredients. Dose, tablespoonful three times a day.

# Da Costa's Solution of Quinino for Hypedermic Use.

R.	Quiniæ sulph	gr. xl
	Sol. acid. tart. sat	m xlviij
	Aquæq s	3 ij
1	1.—Dose according to requirements.	•

# Hughos' Rhubarb Mixture,

'(For Dispepsia.)

R.	Sodii bicarb	3	v
	Tr. nux vom	Z iii	y Z vi
	Tr. capsici	Žiį́	3 ij
	Tr. rhei	Ž	XX.
	Tr. gentian comp	3	xx
7	One to two teaspoonfuls after mea	ls.	

# FORMULARY

OF THE

# NEW YORK MOSPITAL.

(NEW YORK CITY.)

#### FOR EXTERNAL USE.

# **Antisoptic Solutions.**

Sol.	carbolic	acio	l		 ••••	I-20	water
"	44	"			 • • • • •	1-30	"
"	"	"				1-40	"
"	boracic	"			 	1-30	"
"	thymol.		•••••	•••••	 	1-1000	

Distilled water is preferred; it will make a clearer solution than ordinary water.

### White Wash.

R	Potassii sulphuret	
-	Zinci sulphat	3 j
	Aquæ	
	Dissolve each in two ounces of water	and mix,

#### Carbelle Throat Spray. Sodii bicarb..... 3 j Acidi carbolici..... gr. xl Glycerinæ..... 3 vij 3 viij Aquæ.....ad... M. Red Wash. Zinci sulphat..... Spt. lavand. comp ...... Aquæ..... Cochineal coloring..... M. Ward Gargle. Tannin..... Sol. potassii chlorat. sat..... M. Muriate of Ammonia Wash. Ammonii chloridi..... 3 j O ij Tinct. opii..... Aquæ.....ad... Lead and Opium Wash. Liquor plumbi subacet..... 3 iij Aquæ..... Οj ž iv

Tr. opii.....

M.

#### OFFICIAL FORMULÆ

	Alkaline Tar Water.	
Ŗ.	Picis liquidæPotassæ causticæAquæ	ij j v
N	<b>1</b> .	_
-		
	Compound Tincture of Green	Seap.
Tal	ce of—	
	Oil of cade	equal parts
1	И.	•
	Churchill's Tincture of le	line.
R	Iodinii	3 j 3 ij
	Alcoholis	f <b>Z</b> ij
	iedeform Cylinders.	
Ŗ	Iodoformi Tragacanth Mucilag. acaciæ	<b>д ij ss</b> gr. <b>xv</b> q. s.
]	Divide into 10 cylinders, 1 1/2 in. long.	
	Parasiticide.	
Ŗ	Acidi carbolUng. hydrarg. nitrat	gr. x
	Sulphur. precip	3 j
	_	

# **Epilating Stick.**

Ta	ke of—	
1	WaxShellacRosin Burgundy pitchDamar Melt together and roll into sticks of differ	3 iij 3 iv 3 vj 3 x 3 xij rent diameters.
	Coloriess Evaporating Lo	tion.
Ŗ.	Ammon. hydrochlor	gr. xij M. xxxiv <b>3</b> j
	Lotion of Calamino and Zinc	Oxide.
Ŗ.	Pulv. calamin. prep	3 ^j 3 ^{ij} 3 iv
	-	
	Stimulating Lotion.	
Ŗ.	Arnicæ tinct	m xx m xv Z j

#### OINTMENTS.

	Carbolized Cosmoline (Satura	te	d.)	
R	Cosmolinæ	NONO	xx j	
1	Melt each separately and mix.			
0	Intment of Chrysophanic Acid,	Co	ncer	7-
	trated.			
R	Acid, chrysophanic	HOH?	j	
1	Melt the ointment, and while hot add the a dissolved.			in
	Brown Ointment.			
R	Pulv. acid. salicylici	gr.	xl	
	Bals. Peruvian	3	j	
	Cosmolinæ	3	j	
	Ointment of Salicylic Acid	1.		
R	Pulv. acid. salicylici	374	j	
	Cosmolinæ	3	j	
	Ointment of Iodoform.			
R	Iodoformi	1030	j	
	Reduce the iodoform to powder and add he: heat by water-bath till dissolved.	to	the v	as
TAL	ic, near by water battle till dissorred.			

	Cintment of Peruvian Bai	sem.
₽ I	Bals, Peru	3 ij 3 j
·r 1	Dusting Powder.	
ı aı	ke of— Camphor Talc	dr. 1
	Zinc oxideof each	dr. 6
	Compound Oil of Cade Oint	ment.
B	Ol. cadini	f3 j
	Compound lodoferm Cintr	nent.
R.	Pulv. iodoform	3 j 3 j
	Olatment of Tar and Oxide of	f Zinc.
Ŗ.	Ung. picis	3 iv 3 j 3 jss

	Load and Zine Cintme	ent.
Ŗ	Plumbi acetatZinci oxidi	gr. x
	Hydrarg. chlor. mitis	gr. xx
	Olei palmæ purific	3 ss
R	Ointment of Morcury and	lodi <b>ne</b> .
D.	Ung. iodin. comp	<b>3</b> j
_	Tannic Acid Ointment. (Str	onger.)
R.	Acidi tannici	3 j 3 j
	Eczoma Drying Salv	<b>0.</b>
Ŗ.	Plumbi glycerat Ung. zinci oxidi	3 j 3 j
	MIXTURES.	
En	nuision of Cod-liver Oil with Saccharated Lime.	Solution
	(75 per cent. emulsion.)	
Ŗ.	Ol. morrhuæOl. anisiOl. sassafras	3 vj 3 ss gtt. x
•	Liquor. calc. sacchar	3 ij
	Not compatible with acids.	

#### Chlorate of Potessium Mixture.

R	Ammon. muriat		
	Potas«. chloratāā	3	i
	Ext. glycyr	3	SS
	Aquæ cinnamad	ž	iv
1	M.—Dose, a tablespoonful.	•	

# Mixture of ledide of Potessium and Heffman's Anodyne.

•	Potass. iodidi	Sussing	iij j iij
M	I.—Dose, a teaspoonful.		

## Ether Cough Mixture.

R.	Ether. sulph	3	ii
	Tinct. hyoscyam	U	
	Syr. pruni virg		
	Syr. tolutan	3	i
	Aquæad	3	ív
M	Dose, two to four drachms.	Ū	

## Chleroform Cough Mixture.

R	Morphiæ acet	gr. ii
	Tr. belladonæ	3 iv
	Spt. chloroformi	ž vj
	Syr. senegæ	3 j
	Syr. pruni virgad	3 iv
1	M.	•

	Hydrocyanie Mixture	<b>)</b> _	
Ŗ.	Potassii cyanidiSyrupi tolut	gr.	ij
	Liq. morph. sulph., U. S.Pāā	3	i
1	M.—Dose, a teaspoonful.		
	Ward Cough Mixture	١.	
R	Fld. ext. pruni virg	_	iij viij ss x x xviij
	xture of Sulphate of Magnes $(A_{ullet})$	le en	d Ir
<b>R</b> .	Magnes. sulph	97. f. 55.	j xvj ij j <b>vi</b> j
	( <b>B.</b> )		
B.	Magnes. sulph	NO KO KONOMO	j j ij j iij

	( <i>C</i> .)		
Ŗ.	Ferri sulph	gr. 133	xvj j ss j
	Mixture of Iron and Cinche	ona.	
B.	Ferri et ammon. citratis	3 3 3	j ij iv
	M.—Dose, a teaspoonful.		
<b>M</b>	Alkaline Mixture.  ( 4.)  Potass, acetat  Potass. et sodii tartrat  Syr. zingiberis  Aquæ	rangrono	ij j j iij
	_ ( <b>B.</b> )	-	
ĸ	Potass. citratSyr. limonis	3	j ss
1	Aquæ	3	iij
	Nitrous Acid Mixture.		
₽.	Tr. opii deod	13 13 13	ij ss iv

FII:

<b>M</b> i	xture of Morcury and ledide of F $(oldsymbol{A_{oldsymbol{\circ}}})$	Pota	188
R	Hydrarg, bichlor	gr. 3	
N	Tr. gentianāā  M.—Dose, one drachm.	3	j
B.	(B.)  Hydrarg. bichlor	gr. 3	j ¼ iij ij
B.	Hydrarg, biniod	gr. 33	
Ŗ.	(D _• )  Hydrarg, bichloridi  Potass, iodid  Ferri et ammon. cit	gr. 3	j iv j
Ŗ	i Tinct. nuci. vom	3	ij iij
"S	Iix No. 1 and No. 2 at time of dispensi hake Well."	ng, s	and
Ι	Dose, zj., t. i. d.		

# $(E_{\bullet})$

R	Hydrarg, biniod	gr. j
	Potass. iodidi	Ζv
	Syr surant cort	f∄ ij
	Tr. card. comp	fžij
	Tr. card. comp	i <b>∄</b> jv
1	м.	3.

#### Arsenieus Acid Tenic.

R Acidi arseniosi	gr. 1-5
Ferri et quin. cit	gr. lxxx
Tr. cinch. compad	Ĭžij
М	0,

# Compound Acid Tenio.

Ŗ	Tr. nucis vomicæ	f <b>z</b> ij fz iij
	Tr. cinch. comp	f≹ jss
_	Tr. gent. compad	f Z iij
	I.—Dose, two drachms in water, three	times a da

# Hamilton's Tenic.

R	Strychniæ sulph		viij
-	Cinchonidiæ sulph	3	i ´
	Tr. ferri chlor	¥	vi
	Tinct. zingib	3333	ij
	Glycerinæ	·	•
	Acid. phosphoric. dil	3	xvj
3	M.—Dose, one teaspoonful three times	a day	

	Enorvescing Mixturo	-
	No. 1.	
R	Acidi citrici	
	Ferri et quiniæ cit	3
	Aquæ	Ü
	Syr. limonisāā	f3
2	M.	U
	No. 2.	
B	Potass. bicarb	3 ლი
	Aquæad	ηı
]	M.—One fluid drachm of each in two dra	achms o
to l	be mixed at the time of taking.	
	Mixture of Rhubarb and S	ioda.
B		gr. x
	Sodii bicarb	3 i
	Aq. menth. pipad	3 j f3 ij
1	M.	• •
	Rochoile Salt Mixture	
R	Sodii et potass, tart	gr. c
	Ferri et potass. iart	gr. c
	Aquæ menth. pip	f Z iv
	Aquæad	o j
1	M. '	٠,
	Mixture of Squill, Compo	und.
R.	Ammon. chlor	
	Potass. chlorat	ろろう for ii for ii
	Syr. scillæ comp	(5)
	Syr. tolut	f 7 1
	Liq. ammon. acetad	fž v fž ii
	and million access sessions after	

OSBITALS.	125
OF AMERICAN HOSPITALS.	
R Quiniæ sulph	ij iv iij
M. mixture.	
	t. xx
Magnes. Carbon	rtt. ij f3 j 33 iij
M. Mixture	<b>).</b>
M.  Anti-Rheumatic Mixture  (Mistura Antiarthritica.)  R. Potassii iodidi	STATES SS j SS
La Fayette	•
Bals. copai væ	3 ∷
747.	

	Mixture of Digitalis Comp	oun	-
R	Inf. digital	3	≥
-,-	Potass. brom	8	₹.
1	M.	•	<b>)</b>
	*********		
	Compound Infusion of Bu	chu.	,
R	Fol. buchu		
-~	Tritici repentis	3	i
	Potass. acet	7 7	j j
	Acaciæ gran	NOW NOW O	i
	Aquæ fervq. sad	Ö	ίį
1	M.—Ft. infusion.		•
·			
	Mixture of Bicyanide of Mo	reur	у.
R.	Hydrarg. bicyanidi	gr.	SS
	Potass. iodid	3	ij
	Tr. cardam. comp	3	SS
	Syr. sarsap. comp		
_	_Aqueāāad	3	rj
1	M		
	<del></del> 0 <del></del>		
	MISCELLANEOUS.		
			_
	Syrup of Hypophosphites Cer	npou	ınd.
Ŗ.	Calcii hypophos		
	Sodii hypophosāā	gr.	ij
	Potassii hypophos		_
	Ferri hypophosāā	gr.	j
	Acidi hypophos solutq. s		
	Glycerinse	_	
1	Aquæ	3	J
10	<b>1.</b>		

	Bitter Wine of Iron.	
Ŗ.	Ferri et quiniæ cit	gr. lxiv f z ij f z j f z ij f z iv
	М.	•
	Errhine Powder.	
Ŗ.		3 ss 3 ij gr. x 3 ij
	· •	
R.	Fol. sennæ	p.
ΙX	Quassiæāā Potass. bitart Semin. anisi	3 ij 3 j 3 ss
	м.	
	Suppositories of Ergot	
R	Ext. ergot. aquos. (Squibb's) Ol. theobromæ M.—Div. in supposit. No. xij.	Э ij 3 j
	Solution of Acetato of Ammo	nium.
Ŗ		₹ j ₹ xv

## Concentrated Solution of Acetate of Ammonium.

R	Acid. acetic			Mora ij	
				3 ij	
	Ammonii carl	onat		. q. s.	
F	t, sol. neutral.	Evaporate to	3 ij.	This keep	s well.

## HYPORDERMIC SOLUTIONS,

#### Carbolized Distilled Water.

R	. Acidi	carbolici	I part
	Aquæ	destillatæ	999 parts

# Ext. Ergot Solution.

R	Ext. ergot (Squibb's)	I part
	Aquæ destil. carbol	5 parts

# Magendie's Sol. Morphia.

R	Morphiæ sulph	f. 3	lxxx
1	M and filter		

# Hypodermic Solution of Quinine.

R	Quiniæ sulph	grs. clx
	Acid. hydrobrom, (Squibb's)	f. 3 j
	Spts. frumentiad	f. 3 j
7	M. ft. solut.	23.3

# Sol. Pilocarpia Muriato.

Ŗ	Pilocarpiæ	mur	gr. j
	Aquæ dest	carbol	m. j
M	I.—Dose, te		•

# Sol. Apomorphia Muriato.

R	Apomorphiæ muriat. cryst	gr. j M ixxx
M T	I.—Dose, m, x, for emetic. To be prepared only at the time it is wa	nted.

#### POWDERS.

The following powders are sent to the wards and dispensed in bulk, and measured out to the patient in a small measure equal to about 20 grains.

# Pulvis S. I. C.

Ŗ.	Sodii bicarb	600	parts
	Ipecac	1	part
	Cubebæ	300	parts
1	1.—Dose, one measure.		_

### Puivis P. B. S.

R.	PepsinæBismuth		
	Sodii bicarb	100	parts
1	I Dose one measure		•

	Pulvis B. I. C. S.	
Ŗ.	Ipecac	200 part: 3½ " 200 " 400 "
1	M.—Dose, one measure.	
	PILLS.	
	Triplex Pilis.	
R.	Hydrarg, mass	gr. ij gr. j
	Laxative Pills.	
₽ !	Podophil res	gr. ½ gr. ½ gr. ½
	Compound Podophyllin	Pills.
Ŗ.	Res. podophylli	gr. ½ gr. ⅓ gr. j gtt. 1-6
	Rhubarb and Soda Pi	lle.
R.	Rhei pulv	gr. jss gr. 1-10

#### Cormoit's Pilis. Res. podophylli..... gr. 🔏 Ext. nuc. vom..... gr. 1-6 Ext. hyoscyami..... gr. 🔏 M. ft. pil. No. j. Fothergtil's Pills. Morph. mur..... gr. 1-6 Atropiæ sulph..... gr. 160 Pulv. capsici..... gr. 1/2 Pil. aloes et myrrhæ..... gr. j ss M. ft. pil. No. j. Clark's Pills. gr. 3 1/3 gr. j Ouinize sulph..... Pulv. capsici..... Pulv. opii..... gr. 1/2 M. ft. pil No. j. Pills of Lead and Opium. Plumbi acet..... gr. ij Pulv. opii..... gr. i M. ft. pil. No. j. Diuretic Pilis. Pulv. scillæ..... Pulv. digitalis..... gr. j

M. ft. pil. No. j.

Diada 2 Line.
Ferri sulphPotass. carbāā
1. ft. pil. No. j.
Pills of Aloos and Iron.
Ferri sulph. exsic  Pulv. aloes pnrif  Pulv. aromat  Conf. rosæ  M. fiant pil. No. xl.
Antiporiedic Pilis.
Quin. sulph Pulv. capsici Pulv. zingib M. div. in pil. No. xxx.
Pills of Hydrarg. Celecynth, an
Ext. colocynth comp
Pills of Nux Vomica, comp
Pulv. rhei

#### OF AMERICAN HOS

	Compouna Merci
R	Mass. hydrarg
**	Ferri sulph. exsic
	M. ft. pil. No. j.
	Compound Lupu
B	Lupulinæ
	Ext. hyoscyami
	Pulv. acaciæ
	M. ft. pil. No. j.
	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
	Compound Pills of I
R	Ext. hyoscyami
	Heydrarg. proto-iod
-	Ferri et quin. cit
1	M. et ft. pil. No. j.
	Compound lodol
R	Iodoformi
-	Hydrarg, chlorid, corros
	Ferri redacti
-	M. ft. pil. No. j.
	_
	Compound Chino
R	
IX	Chinoidinæ
	Ferri redacti
1	M. ft. pil. No. j.
4	at the pile 140. J.

# ROOSEVELT MOSPITAL FORMULAR

(NEW YORK CITY.)

#### TONICS.

	Cod-Livor Oil Emulsion.		
R.	Cod-liver oil	3333	ij j j ss
	God-Liver Oil with Hypophos	اط	tes
B.		gr.	x
	Hypophos. soda	gr.	V
	Hypophos. potass	gr.	¥
	Hypophos. iron  Dil. phosphoric acid, s. q	gr.	V
	Cod-liver oil	3	ij
	Glycerine	3	J
	Boiling water s. q	3	SS
1	M.—Dose, 3 ss.		

#### OF AMERICAN HOSPITALS.

#### Arsonicai Mixturo.

R.	Sol. potass. arsenite	Μj
-	Comp. infus. gentian	3 ss
	Water, s. q	Žј
N	1.—Dose, z j.	

#### Iton and Arsonic.

R	Sol. chlorid: arsen	ηij
	Water, s. q	3 j
1	M.—Dose, z i.	

# Compound Cod-Liver Oil.

R	Cod-liver oil	<b>3</b> ss
	Syr. lactophosph. lime	Ψ̈́x
	Syr. iodide iron	m iv
	Sol. lime, s. q	3 j
1	M.—Dose, z i.	-

# Laxativo Iron Mixture.

R.	Sulph. magnesia	<b>3</b> 88
	Sulph. iron	grs. ii
	Dil. sulph. acids. q	
	Comp. tinct. gentian	m xv
	Water, s. q	m xv 3 ss
	M.—Dose, Z ss.	_

A. solution.

#### Child's Tonic. Ammon. cit. iron..... grs. v. Sol. arsenite potass..... m, ii 3 88 Syr. orange..... Žј Water, s. q..... M.—Dose, Z i for a child three years old. **Quining, Iron and Arsenic.** R. Sol. arsenit. potass..... m, iij Cit. iron and quinine..... gr. v Comp. tinct. gentian s.q..... 3 ij M.-Dose, 3 ij Bright's Tonic. Sulph. strychnine ..... gr. 1-32 Tinct. chlor. iron..... m xv Simple syrup..... 3 j 3 ij Water, s. q..... M.-Dose, 3 ij Effarvoscing iron. Cit. iron and quin..... grs. 5 1/8 Citric acid..... grs. 7 1/2 Water, s. q..... 3 ij M.-Label, A. solution. Dose, 3 ij; mix with 3 ij B. solution. Bicarb potass..... grs. 1834 Water, s. q..... 3 ij M.—Label, B. solution. Dose, Z ij; mix with Z ij

OF AMERICAN HOSPITALS	OF	AMER	ICAN	HOSPITAI	s.
-----------------------	----	------	------	----------	----

# Tincture of iron.

<b>₽</b>	Tinct. chlor. iron	m xv
	Glycerine	<b>3</b> ss
	Cinnamon water, s>q	ž ij
1	1.—Dose, z ii.	-

# COUGH MIXTURES.

# Mixture No. 1.

Ŗ.	Sulph. morphine	gr. 1-16
	Chloride ammon	gr. 21/2
	Syrup tolu	m, xv
	Water, s. q. to	<b>3</b> ij
1	M.—Dose, 3 ij.	•

# Mixture No. 2.

R Fl. ex. wild cherry bark	m 55%
Sol. cyanid. potass. { 16 grs. } uni	m 15-16
Simple syrup, s. q. to	<b>3</b> ij
M.—Dose, 3 ij.	

# Cedoino Mixturo.

Ŗ.	Codeine Dil. phosphoric acid	gr. 1-6 m. v
	Simple syrup, s. q. to	βj
Ŋ	1.—Dose, z j.	

	ipocae Mixturo.	
R	Wine of ipecac Muriat. of ammon Camphor water, s. q. to	m v gr. v Ž ss
1	M.—Dose, 3 ss.	Ü
	Antimony Mixturo.	
B.	Tartrat. antim and potass	gr. 1-24 gr. 1-10 3 j
	Sanguinaria Mixture	_
R		m v grs. v 3 j
М	Dose, 3 j.	•
	Carbon. and Iodido Ammon. I	Mixture.
Ŗ.	Carbon. ammon	grs. iij grs. ij 3 j
1	M.—Dose, 3 j.	
	Atropino Mixture.	
Ŗ.	Sulph. atropine	gr I-128 grs. iij
1	Simp. syrup, s. q. to	<b>3</b> j
	, a ,	

	Childron's Mixture.		
Ŗ	Paregoric	m,	x
7-	Muriate ammon	grs.	
	Syrup squills	ďη	x
	Syrup tolu, s. q. to	3	j
	M.—Dose, 3 j.	_	
	Hydrobromic Mixture		
R	Dil. hydrobromic acid (10 per ct.)		
	Spirits chloroform of each	m.	iv
	Syrup squills	m.	viij
	Paregoric, s. q. to	3	
	Bromide and Hyescyamus N	lixtu	Ire.
Ŗ.	Bromide potass	grs.	ХX
	Tinct. hyoscyam	3	SS
	Camphor mixt., s. q. to	3	SS
	M.—Dose, 3 ss.		
	Squilis Mixture.		
R	<u> </u>	100	
ΙX	Tinct. squills Wine of ipecac	m	
	Syrup of tolu		iv xx
	Water, s. q	3	
	M.—Dose, 3 j for child one year old.	Э	,
	——————————————————————————————————————		
	Chloride of Ammen. Mixt	ure.	
R	Chloride ammon	grs.	iı
	Syrup wild cherry bark	_	58
	Water, s. q. to	3	j
	M.—Dose, 3 j for child one year old.	••	•

B.	Emphysema Mixturo.  Iodid. potass	gr. v m v m xlv m ij ss 3 ss
	DIARRHŒA MIXTURE	<b>S.</b>
R.	Nitric acid.  Nitric acid.  Tinct. opium.  Tinct. camphor.  Comp. syr. rhubarb.  Syr. ginger.  Water, s. q.	m xx m vijss m vijss 3 j 3 j
<b>R</b>	Kino and Opium (Dysenter Tinct. kino	ory.) m xx m xx m xv m v
	Bismuth and Soda.	
R.	Sub. nit. bismuth	grs. iij grs. ij 3 ss 3 ij

M.—Dose, 3 ij.

•	Oplum and acid.	
R.	Comp. sol. opium, Squibb's Arom. sulph. acid	m x m x
	Company Diameth	
B.	Campound Bismuth.  Camph. tinct. opium  Subnit. bismuth  Chalk mixture, s q  M.—Dose, 3 ij as directed. Shake.	m x grs. x 3 ij
	Compound Chalk.	
R.	Camph. tinct. opium  Tinct. krameria  Comp. spirits lavend  Chalk mixture, s. q  M.—Dose, 3 j, as directed.	m v m v m ij 3 j
	Arematic Rhubarb.	
Ŗ.	Powd. bicarb. soda	grs. ij m xx gtt. v 3 ss 3 j
I	M.—Dose, 3 j.	
	Chalk and Catechu.	
B.	Tinct. opium	m vijss 3 j 3 ij

# OFFICIAL FORMULÆ

	Colo.		
R.	Fl. ex. coto bark	m 3 3	ij SF j
Ŗ.	Castor oil. Powd. acacia. Sugar Tinct. opium. Spirits nutmeg. Orange flower water.	m	ss iv xx
1	M.—Dose, 3 j.		
	MIXTURES FOR DYSPEPSIA R. & S. Compound.	1, E1	rc.
R N	Tinct. nux vomica	gr. grs. grs.	¼ v ij
Ŗ.	Powd. rhubarb	grs. grs. gr. 3	v X

	Pepsine.	
R	Saccharat. pepsine Dil. muriat. acid	grs. iij M iij
	Glycerine	m, v
	Water, s. q	a j
	M.—Dose, 3 j.	•
	Dyspepsia Mixture.	
Ŗ.	Dil. hydrochloric acid	m, x
	Tinct. nux vomica	ηx
	Tinct. cinchona, s. q	3 j
	M.—Dose, 3 j.	
	Celic Mixture.	
R	Spirit of chloroform	m, xx
	Arom. spirit of ammon	m xx
	Comp. spirit lavend	m xx
	M.—Dose, 3 j.	
	infant Laxative Mixture	<b>).</b>
R	Nitrate potass	gr. j
	Sulph. magnes	grs. v
	Lemon syrup	3 8s
	Water, s. q	ž i
	M.—Dose, 3 j for a child one year old.	., ,
	Rhubarb and Magnesia Mix	ture.
Ŗ		grs. v
	Tinct. rhubarb	m, iii
	Syr. ginger	m, x
	Peppermint water, s. q	ã j
	M.—Dose, 3 j for a child one year old.	-

#### Caster Oii.

Ŗ.	Castor oil	3	j
	Glycerine	3	SS
	Ol. cinnamon	m	1/8
ľ	M.—Dose, 3 jss.		

# Vinum Gastricum.

Take one perfectly fresh calf's rennet, cut up into pieces and put in a jar, and pour on one quart of good sherry; do not filter. Dispense supernatant fluid after standing thirty-six hours.

#### SEDATIVES.

# Chioral.

Ŗ.	Chloral hydrateBromide potass		vij ss vij ss
	Orange flower water	m	•
	Water, s. q	3	j
N	I.—Dose, z j.		

#### Jim Jam.

Ŗ.	Bromide potass		xxx.
	Magend. sol morph	grs. m	
	Syr. orange peel		
	Water, s. q	ž	j i <b>v</b>
1	M.—Dose. Z ss. as directed.		

# Mixed Bromido.

Bromide sodium Bromide potass Bromide ammon Syr. orange peel Water, s. q M.—Dose, 3 iv.	grs. xx grs. x 3 ij
Bremid	• Solution.
B. Bromide sol	
-	
Hys	storia.
R. Tinct, castor	s. q 3 j
Sound	
Tr. aconite root Sulph. morph Water, s. q M.—Dose, 3 ij, as dire	gr. 3-16 3 ij
- Elixir o	 f Quarana.
R Fl. ex. guarana Simple elixir. s. q	

# DIURETICS.

	Buchu Mixture.		
R.	Fl. ex. buchu	3	SS
	Tinct. hyoscyam	ž	22
	Sol. citrat. potass	3	ss ss j
I	M.—Dose, 3 ij.	Ü	•
	Digitalis.		
R	Infus. digitalis	3	i
	Acet. potass	grs.	
	Water, s. q	3	
1	M.—Dose, g ij.	•	•
	ANTI-SYPHILITIC MIXTU	RES	<b>.</b>
	Wixed Treatment.		
Ŗ.	Bichlorid mercury	OT.	1-3
~	Iodide potassium	grs.	
	Comp. syr. sarsap		
	Water, s. q	2	ss ij
,		o	4
1	M.—Dose, 3 ij.		
	lodide Potassium.		
R	Lodide potassium	grs.	x
	Comp syr. sarsap	•	IV
	Water, s. q	3	
1	M.—Dose, 3 j.	ð	•
	Setunded Cal Ladds Date		_
	Soturated Sel. Iodide Potas	suu	۳,

 ${\mathfrak Z}$  j sol. represents  ${\mathfrak Z}$  j iodid. potass.

OF AMBRICAGO HOSP

#### **MUSICELLA MED**

	liadide and Gold
R	Tofide potasiium
1	Wine enlichteum seed
	Сапр. гутар запар
	Water, s. C.
3	L.—Duse, I sa.
	Ont Ontandate
	Sol. Salicylate
R	Salicylate soda
	Salicylate soda. Bicarlı soda, y. q. Powd. est. licteire.
	POWE CAL BETWEE
	Glycerine
	Water, s. q.
3	L.—Dose, 3 ij.
	The second second
	Infant Febrifi
R	
B.	Citrat. potass
B	Citrat. potass.  Bicarb. potass.  Syr. lemon.
R	Citrat. potass.  Bicarb. potass.  Syr. lemon.
	Citrat. potass
	Citrat. potass.  Bicarb. potass.  Syr. lemon.
	Citrat. potass. Bicarb. potass. Syr. lemon. Water, s. q I.—Dose, 3 j for child 1 year
2	Citrat. potass.  Bicarb. potass.  Syr. lemon.  Water, s. q
2	Citrat. potass.  Bicarb. potass.  Syr. lemon.  Water, s. q
2	Citrat. potass.  Bicarb. potass.  Syr. lemon.  Water, s. q.  6.—Dose, 3 j for child 1 year  Lafayette Min Copaiba.  Spt. nitrous ether.
2	Citrat. potass. Bicarb. potass. Syr. lemon. Water, s. q. Lafayette Mb Copaiba Spt. nitrous ether. Comp. spt. lavender
2	Citrat. potass. Bacarh. potass. Syr. lemon. Water, s. q
R.	Citrat. potass. Bicarb. potass. Syr. lemon. Water, s. q. Lafayette Mb Copaiba Spt. nitrous ether. Comp. spt. lavender

R.	H. Fermula for Sol. Merph. Ma Hypodormic Use.	gend.	for
Ŗ.	Salicylic acid  Distilled water, s. q	grs. xvj gr. ½ 3 j	
	Dissolve the salts in the water previously hing point. Filter.	neated to	the
	Saturated Sol. Chierat. Pote	)SS.	
	<b>1—16</b> .		
	<del></del>		
	Sol. Erget for Hypódormie (		
Ŗ.	Ext. ergot, Squibb's	3 ij 3 ss 3 jss	
N	<b>I</b> .	0.	
	<del></del> .		
	Ward Solut. Atropino.		
Ŗ.	Sulph. atropine	gr. 1-96	
N	Mater	<b>3</b> J	
	Chloroform and Whisky.		
Ŗ.	ChloroformWhisky, s. q	m x	
N	M.—Dose, 3 ss.	0 = 3	

#### PILLS.

	Compound Arsenic Pilis	B.
R.	Arsenious acid	gr. 1-20 gr. j grs. ij
	Aloe and Iron.	
R.	Dried sulph. iron	gr. j grs. j ss
	R. S. and i. Piil.	
R.	Powd. rhubarb	gr. j gr. j gr. 1/8
	lodoform.	
	IodoformMake one pill.	I gr
	Company Diversity	
B.	Corrosive sublim	gr. 1-32 gr. j gr. j

Н	yoseyamus and Oxide Zinc.
R N	Ext, hyoscyamus
	Quinine (1 grain).
Ŗ.	Sulph. quinine
	Honey, s. q
Ŋ	M.—Make one pill.
	-
	Quinino (3 greins.)
B.	Sulph. quin
	Honey, s. q
N	1.—Make one pill.
	Biand's.
R	Sulph. iron
	Carb. potass
	Pulv. tragacanth, s. q
N.	I.—Make one pill.
	Laxative.
R.	Ext. nux vom
	•

OF A	MERI	CAN	HOSP	ITALS.
------	------	-----	------	--------

151

#### Codeine.

# CAPSULES.

#### **POWDERS**

# Bismuth, Pepsine, and Morphine.

10 grs. 1-6 gr.

#### Sentenine

# Infant Calomei.

R. Calomel......gr. j Sugar.....grs. iij

M.-Make one powder.

Ŧ	

#### OFFICIAL FORMULÆ

# Ciark's Powder.

Ŗ	Sulph. quin	grs. x grs. v gr. j
N	I.—Make one powder.	8 1

### POWDERS IN BULK.

Dose, one measure represents about twenty grains.

# Calemel and Jaiap Comp.

Ŗ.	Calomel	grs.	x	
	Comp. powd. jalap	grs.	x	
7	<b>v</b> f.			

# Sode.

R	Bicarb.	soda	grs.	хx
TX	Dicai D.	30ua	K12.	**

# Bismuth and Sada.

Ŗ	Subcarbon, bismuthBicarb, soda	grs. grs.	
M	ſ.	Ū	

# ipecac and Soda.

Ŗ	Pulv. ipecac	gr	*	
	Bicarb. soda	ørs.	хx	

M.

#### OF AMERICAN HOSPI

	Bismuth, Cubeb, ar
R	Subnit. bismuth
	Pulv. cubeb
	Pulv. bicarb. soda
	M.
	The same of the sa
	Bismuth and Pe
R	Subnit. bismuth
	Pepsine
	M.
	The second second second
	Alkaline Laxato
R	Chloride sodium
	Bicarb. soda
	Carbon calcium
	Sulph. soda
	M.—Dose, 3 ij.
	EXTERNAL APPLICA
	Evaporating Loti
R	Muriat. ammon
1,x	Alcohol
	Water, s. q
	M.—External use.
	Red Wash.
R	Sulph. zinc
**	Comp. spirit lavend
	Water, s. q
	M.

	Churchili's Tinet. ledine.
R	Iodine grs. lxxv Iodide potassium grs. xv
	M.—For external use.
	Sol. Carbolic Acid and Glycerine.
-,-	Carbolic acid, pure
teı	M.—C. A. and G. As a base for dispensing for in- rnal use.
	Comp. Tinct. Green Soap.
Ŗ	<b>=</b>
of	Dissolve the green soap in the oil of cade with the aid heat. Add the alcohol and shake. For external use.
	Sol. Load and Opium.
Ŗ	Sol. cubacet. lead
	M —External use.
	Alkaline Lotion.
Ŗ	
	Biborat soda grs. x Cherry laurel water
	M.

# Sulphurous Acid Lotion. Sulphurous acid..... Water..... M.—External use. Cemp. Chioroform Linimont. Chlorof. commercial.....) Tinct. capsic..... M -External use. Liniment Stimulant. Ammonia water..... Tinct. capsicum..... Soap liniment..... M.—External use. **Boratod Sciution.** Salicylic acid..... I part. Boric acid..... 6 parts. Water..... 500 M. Chloreform and Acenite Liniment. R. Commercial chloroform..... Tinct. aconite root..... Soap liniment.....

M.—External use.

# OFFICIAL FORMULA

	Ward Gargie.		
Ŗ	Biborate soda	3	j
	Sat. sol. chlorate potass	3	j
	Water, s. q	3	V
1	M.—Use as a gargle.		
	Chioride Zinc Gargio.		
Ŗ.	Chloride zinc	grs.	x
	Glycerine	3	88
	Water, s. q	3	iv
I	M.—Use as a gargle.	_	
	Tannin and Alum Gargi	· •	
R.	Alum		SS
•	Tannin	NO ROMON	SS
	Glycerine	3	SS
	Water, s. q	3	iv
1	M.—Use as a gargle.		
	Aikaiino Tar Water.		
R	Liquid tar	Z	ii
	Caustic potass	美	i
	Water	ž	ij j V
]	M.—External use.	Ū	
	OINTWENTS.		
	Salicylic and Zinc Ointm	ont.	
Ŗ	Salicylic acid	gr	s. xv
	Powd. oxide zinc	- {	ζijs ζijs
	Powd. starch	ž	z ijs
	Petroleum	ž	5 j
	М.		

	Carbolic Olatment, 3 por	ct.	
R	Carbolic acid	gr.	I
	Petrolatum		331/3
	-	P- 2-	33/3
	· Cade and Zinc.		
R	Oil of cade	Z	i
	Oxide of zinc ointment	3	í
7	đ.	J	,
-			
	i <b>o</b> doform.		
R	Iodoform	3	i
	Petrolatum	3	í
7	Л.	9	,
-			
_			
Re	isam Poru Ointment (Strong E	lraw:	n Mint
	. Bus i ara amman (anona r		
	ment.		
	Salicylic acid		part
	Salicylic acid	1	
	Salicylic acid	1	part
Ŗ.	Salicylic acid	1 : 5 :	part parts
Ŗ.	Ment. Salicylic acid	1 : 5 :	part parts
Ŗ.	Ment. Salicylic acid	1 ; 5 ] .11	part parts
Ŗ.	Salicylic acid	. 11 . 11	part parts "
R.	Salicylic acid	. II . II 	part parts "
R M R	Salicylic acid	. II . II 	part parts "
R M R	Salicylic acid	. II . II 	part parts "
R N R N	Salicylic acid	5 .11 (iid).	part parts " part
R M R	Salicylic acid	5 . 11	part parts  part part part part
R N R N	Salicylic acid	5 . 11	part parts " part

#### **Tube lodoform Cintment.**

R	Expressed oil almonds	9 parts
	Spermaceti	I part
	Wax	ı ""

Melt together at a moderate heat the oil of spermaceti and wax, stirring the mixture constantly until it is nearly cool, then add powdered iodoform in the proportion of one part of iodoform to four of the ointment, and continue the stirring until it has become uniformly soft.

# Ohrysarobin Golatine.

Dissolve one part pure gelatine in two parts water in water-bath, and add ten per cent. chrysarobin.

# Pyrogalioi Gelatine.

10-20 per cent.

# Salicylated Gelatine.

5-10 per cent.

# **Dusting Powder.**

R	Camphor	3	j Vi
	Oxide zinc	3	νj
1	٧٢.	_	

# DEPARTMENT OF

# Public Charities and Correction,

' (NEW YORK CITY.)

# BELLEVUE HOSPITAL.

# Hydrocyanic Mixture.

B	Potass. cyanidi
-	Morphiæ sulphatāā gts. 4
	Syr. tolut fl. 3 4
N	M.—Each 3 contains 1/8 grs. each of potassium cya
nid	e and morphia sulphate. Dose, a teaspoonful.

#### Rhubarb and Soda.

R	Sodii bicarbon		3	I
	Pulv. rhei		¥	1/2
	Pulv. rhei	fl.	ž	2
	Aquæ,q. sad	fl.	3	4
N	1.—Dose, a tablespoonful.		Ū	•

# Tincture of Phospherus.

withing up 1	Phosphori
	Anti-Rhoumatic Mixture.
R	Sodii et potass. tart

# Thomson's Tonic.

M.—Dose, a teaspoonful.

R	Ferri et ammon. cit		1 30
	Tr. gentian comp	fl. 3	2 1 1/4
	Aquæq. sad	fl. 3	1 ½ 8

Dr. W. H. Thomson

#### OF AMERICAN HO

# **Phosphorated Cod** Olei phpsphorati* (1 per cen Ætheris..... Olei morrhuæ.....q. s. M -Two hundred and thirtycally 1/2 fl. 3, contain 1-30 grain phosphorated oil should be weigh OUT-DOOR DEPARTM Whooping-Cough Tinct. nucis vom..... Vin. ipecac..... Syr. sarsap. comp..... " senegæ..... M .- Dose, a teaspooned, for *Oleum Phosphoratum. R Phosphori..... Olei morrhuæ..... This is a r per cent, solution of phosp posed by Dr. E. R. Squibb. It is made contains the full amount of phosphorus. contents is to be used, a few drops of et the vial, before it is again corked and ser form on the surface or at the bottom, the as to leave this in the vial. Should this precipitate make its appearance, a fresh It is best to add the whole coments of cod-liver oil to be ready for administra kept on hand in full and well-closed bott the dark.

When using the phosphorated oil it sh

zoeight:

# OFFICIAL FORMULÆ

Ŗ	Cough Mixture.  Syr. tolut.  Syr. pruni. virg		
	Spt. ætheris comp	Ü	
	Expectorant Mixturo for C	hlidro	m.
R	Syr. senegæ " pruni virg	fl. 3	
	M.—Dose, a teaspoonful	Dr. E	Folgate.
_	Mistura "Bronchi."		
R	Ammonii carbon		10 1½ 1
	Aquæq. sad	fl. 3	2
	M.—Dose. a teaspoonful for children.  Dr. Beve	erly Ro	binson.
	Carboneto of Ammonia M		
R	Syr. senegæ  " ipecac  " tolut  Ext. glycyrrhizæ  Aquæ cinnamq. s. ad	fl. 33 fl. 33 fl. 33	1/2 4 2 4 1/2 4
	M.—Dose, a teaspoonful, for children.  Dr. G.	co. H.	Bosley.

	Chloride of Ammonium M	lxt	ur	D.
R	Ammonii chloridi		3	1/2
	Potassii chlorat	و	rs.	4Ó
	Syr. senegæ	fl.		4
	" ipecac	fl.		
	" tolut	fl.	3 3	3 . 5
	Ext. glycyrrhizæ		ž	Ĭ
	Aquae cinnamq. sad	fl.	ž	4
	M.—Dose, a teaspoonful, for children.		·	•
		en.	ς.	Boslev.
			٠	som,
	Expectorant Mixture			
R.	Syr. scillæ comp			
	Syr. ipecac	fl.	3	I
	Syrupi	fl.	3	I
	M.—Dose, a teaspoonful, for children.		U	
	2000, a tomopounar, for omnatour	D	r. S	wezev.
	Licorice Mixturo.			
₽.	Ammonii chloridi			
	Ext. glycyrrhizæāā		3	2
	Tr. opii camph	fl.	333	2
	Aquæq. sad	fl.	3	4
	M.—Dose, a teaspoonful.			
		D	r. 1	Brekes.
_	Sedative Mixture.			
Ŗ	Acid. hydrocyan. dilut	а	_	_
		fl.	3	1
	Tr. hyoscyami			
	Syr. tolutani		7	_
	Aquæ camphoræāā	п,	3	I
	M.—Dose, a teaspoonful.			
	n Dr.	K	atze	mbach.

He	Hmann's Anodyne and lodid	o of	Potash.
R	The second secon		3
	Ext. pruni virg. flāā		1
	Syrupi		1
	Aquæ	fl. 3	ĩ -
1	M.—Dose, a teaspoonful.  Prof. E.	G. J	aneway.
	Nitrate of Potash Mixt	ure.	
R	Potass, nitrat	gr	. 1
	Syr. ipecacāā	fl. 3	1/2
	Syr. pruni virgq. sad	11 3	2
N	1.—Dose, a teaspoonful, for children	9	
	Chlorate of Potash Mix		
R		ture.	1
_	Ext. glycyrrhizæ	3	1/2
	Ammonii chloridi	0 3	1
,	M.—Dose, a teaspoonful.	fl. 3	4
-	Dr. Gco.	G. 1	Vheelock
H	ofiman's Anodyne and lodide	of I	Potash.
R			
	Potass, iodidi Syr. pruni virg	3	13
	Spt. æther. comp	fl. 3	11/2
1	M.—Dose, a teaspoonful.		
	Di	. Kat	senbach.

# OF AMERICAN H

Cyanide and Bromide

B.	Potass. bromidi
	Optum, Rhubarb a
R.	Tr. opii
	Castor Oil
R.	Olei ricini
	and the same
	Rhubarb ar
R	Sodii bicarb

# OFFICIAL FORMULÆ

Bicarbonate of Soda.
R. Sodii bicarb
Rhubarb and Lime.
R. Tr. opii camph
Dr. Ackerman
Pulv. Bismuthi, Greta ot Opii.  3 Bismuth. sub-nit
Pulv. Bismuthi et Deveri.  Bismuthi sub-nit
Pulv. Bismuthi et Pepsini.
Bismuthi sub-nitr

# Pulvis Bismuthi Comp.

£	Bismuthi sub-nit	
	Pulv. sacchari	
	Pulv. acaciæ	
	Pulv. zingiber	
1	M.—Dose, a tablespoonful, for adults, in dyspepsia.	
•	Dr. Wheelock.	
	Salicylic Acid Mixture.	
B	Acidi salicylici grs. 160	
	Potass. acetat grs. 320	
	Glycerinæ fl. 3 1	
	Glycerinæ fl. 3 1 Αςυq. sad fl. 3 4	
1	M.—Dose, a teaspoonful.	
	Syrup of Biniodide of Marcury.	
R	Potassii iodidi grs. 80	
	Hydrarg. biniodidi gr. 1½ Syrupi fl. 3 2	
1	Dissolve and mix. Dose, a teaspoonful.	
-	Dr. Banks.	
	Thompson's Mixed Treatmont.	
R	Hydrarg. biniodidi gr. 1	
	Tr. aurantii fl. z i	
	Potassii iodidi	
1	1.—Dose, a teaspoonful.	
-	• • • • • • • • • • • • • • • • • • •	
	Dr. Beverhout Thompson.	

	•
	Taylor's Mixed Treatment.
$\mathbf{R}$	
	Potass. iodidi 3 4
	Syr. sarsap. comp
	Aquæ
1	M. Dr. R. W. Taylor
	98 - 1-40 - A-1-1 981 - 4
D	Muriatic Acid Mixture.
Ŗ	Acidi muriatici
	Aquæ
7	I.—Dose, a teaspoonful.
	1.—Dose, a teaspooniui.
	Pil. Ferri Quin. ot Strych.
R	Quiniæ sulph
	Ferri redacti
	Strychniæ acetat gr. 1
	Ext. gentian q. s.
N	I.—Divide into sixty pills.
D	Iron and Citrato of Ammonia. Ferri et ammon, cit
ΙX	Ammonnii carbonāā grs. 32
	Syrupigrs. 32
	Aquæanisi
N	I.—Dose, a teaspoonful.
	Dr. J. L. Smith
	iren and Cinchona.
$\mathbf{R}$	Cinchonse sulph 3 1
	Tr. ferri chloridi
_	•
M	.—Dose, a teaspoonful

# OF AMERICAN I

	Lemon T
R	Cinchonæ sulph
	Sulphate of C
R	Cinchonæ sulph
1	1.—Dose, a teaspoonful.
-	and the designation of the same
	Iodide Mi
B	Potassii iodidi Syr. ferri iodidi Tr. calumbæ
0	M.—Dose, a teaspoonful.
Ex	ternal Applications
	Ung. Hydr. Ox. Re
R	Hydrarg, oxidi rubri Plumbi acetat Cerati

	OF AMERICAN
En	nulsion—
	Cod Liver 0
R	Olei morrhuæ Liquor potassæ Mellis, Pulv. acaciæ Ol. anisi Ol. menthiæ vir I.—Dose, a tablespoonful
	_0
	CHARITY HOSPITA
	Expectoran
R	Tr. sanguinar
-	I Dans a terraneous ful

# Asthmatic

R Spt. æther comp....... Liquor morph, sulph. (U M .- Dose, from one teasp

Hydrocyanic Mixture	٠.		
Potass. cyanidi Vini antimonii			
Mucil. acaciq. s. ad	Я. Я.	<b>3</b>	½ 1
1.—Dose, one teaspoonful.			
Hot Drops			
Tr. opii	Я. Я.	33	2 I
Anti-Emetic Minture			
Creosoti		m	) 2 30
Aquæq. s ad  M—Dose, a teaspoonful.	fl.	25.23	2
Anti-Rheumatic Mixtu	r		
Sodi; et potass, tart	fl, fl.	33300	1/2 2 2
	Potass. cyanidi Vini antimonii Syr. tolut Mucil. acaci	Vini antimonii	Potass. cyanidi

lodide of Potash Mixtu	re.	
R Potass, iodid	3	4
Tr. gent. comp	fl. 3	I
Lemon Tonic.		
R. Cinchonæ sulph	fl. 3	a s

OF AMERICAN HOSPITALS.

173

# Solution Ergotine.

Cod-Liver Oil Mixture.

Olei cinnamomi.....

R Olei morrhuæ.....

M.—Dose, a tablespoonful.

R.	ErgotiniGlycerinæ	gr.	36
	Aquæāā	m,	108

M.

# INSANE ASYLUM, BLACKWI

## Expectorant Wixture. R Spt. etheris comp ..... Syr. ipccac..... Tr. opii camph ..... M.—Dose, a teaspoonful. Cannable Mixture. R. Tr. cannabis Ind ..... Spt. menthæ pip..... Aquæ ......q. s. ad... f M.—One dose. To be taken thrice daily **Epileptic Bolladonna Mixt** R. Potass. bromidi..... Tr. belladonnæ ..... Aquæ ......q. s. ad... f M.—One dose to be taken thrice daily. Sedativo Mixture. R Chloralis..... Extr. conii sem. fl..... Aquae .....q. s. ad... f M.—One dose, to be taken thrice daily,

Epileptic (Conlum) Mixture.  R. Potass. bromidi
R. Potass. bromidi
lowing the attack, indication of cerebral congestion an especially where hemmorhage is feared.  Dr. Chas. R. Smith.
Strychnine Tonic.
R. Tr. ferri chloral
NINETY-NINTH STREET RECEPTION HOSPITAL.
Piliula Aloos ot Foliis.
R. Ext. aloes
constipation.

	•	
	Laxative Pilis.	
R.	Pulv. rhei	
	Pulv. aloes	15
	Ext. bellad	•
	Ext. nucis vom	
	Resin. podophylli grs.	2
	Olei caryophylli gtt.	2
	A.—Divide into twelve pills. Dose, one pill m	
and	d evenings.	iornings
	Anti-Epileptic Pill.	
R.	Argenti nitrat gr. 1	ю
~	Zinci oxidi gr. 2	
		q. s
1	M.—Divide into twenty pills. Dose, one pil	
dail		unice
uan	<u></u>	
	Greton Oil Linimont.	
R	Olei tiglii fl. 3	2
	Olei olivæ	-
	Olei terebinthinæ	
	Aquæ ammoniæ	
	Spt camphoræāā p.e. q.s. ad fl. 3	2
3.	M.—S. Externally in chronic muscular pains.	-
	m.—5. Externally in chronic muscular pains.	
	· · · · · · · · · · · · · · · · · · ·	
	HART'S ISLAND HOSPITAL.	
	Anti-Rheumatic Mixture.	
Ŗ.		Ī
	" acetat 3	1
	Tr. colchic sem fl. 3	2
_	Aquæ 0 :	2
M	M.—Dose, a tablespoonful.	

OF AMERICAN HOSPITAL
----------------------

### 177

## **Expecterant Emulsion.**

Ŗ	Morph. sulph	grs. 2
	" ipecac	fl. 🐧 2
	" pruni virg	fl. 3 1½
	Tr. sanguinariæāā Aquæ	fl. 3 ½ fl. 3 2
V	I.—Dose, a teaspoonful.	

### Diarrhea Mixture.

Ŗ.	Tinct. capsici	fl.	3	I
	Catechu			
	" kino	_	_	
	" krameriæ       āā         " opii          Spt. menth. pip	Ħ.	3	4
	" opii	Ħ.	3	3
	Spt. menth. pip	fl.	3	2
	Aquæāā	fl.	3	4
N	1,—Dose, 30-60 minims.			

## INFANT'S HOSPITAL, RANDALL'S ISLAND.

## **Whooping-Cough Mixture**

Ŗ	Acid. nitric. dil	fl. fl.	3333	I 1/2 2
	I _Dose a teaspoonful		•	

# Pulv. Bismuthi et Pepsini.

Ŗ.	Pimuthi sub-carb		
	Pepsiniāa	grs.	2
M	I.—One dose or children.	_	

### Diarrhosa Mixture.

Ŗ	Bismuthi sub-carb	grs.	
	Pulv. ipecac comp	•	_
M.	I.—One dose for children in diarrhœa.		

### Cough Wixture for Infants.

R.	Tr. opii camph		
	Spt. ammon. arom	fl.	ξı
	Ext. ipecac fl	fl. a	* %
	Svr. pruni virg	A. 3	ľ
	Aquæq. s, ad	fl. 3	3
1	M.—Dose, a teaspoonful.	•	, ,

## FORMU

OF TI

# Long Island Co

(BROOKLY

MIXT

## 1.—Mixture of Citrate

R Ferri et quiniæ cit..... Syr. simplicis..... Aquæ...... M.

### 2.-Mixture of P

	3.—Mixture of Iron and Cir	nche	ma.
₽. -	Cinchoniæ sulphatis	BRITA	j iv j i▼
1	M		
	4.—Mixtura of Sulphate of	Quis	ia.
R	Quiniæ sulphatis	grs. q. s.	
1	M.	Ü	•
	5.—Mixture of Acetato ef	lron.	
Ŗ.	Tr. ferri chloridi		ij sa j ss
1	Aqua	3	j
•			
	6.—Mixture of Rhubarb and	Sot	la.
₽. 1	Ext. rhei ¶	m m grs.	iij xl

	7.—Hydrocyanic Gough Mix	ctur	<b>9.</b> ·
Ŗ.	Morphiæ sulphatis	gr. 3	ij ss
M	Syr. tolutanis	3	ij
	<del></del>		
	8.—Bromide Cough Mixture	(Re	nd).
Ŗ	Potass. bromidi	3 Karana	ij ij j iij
	Belladonna Cough Mixtu	re.	
Ŗ.	Ammon. chloridi           Morphiæ sulphatis           Tr. aconiti rad           Ext. belladonnæ fl           Ext. glycyrrhizæ fl           Syr. simplicis         āā           Aquæ         q. s. ad	.m	ij xvj
Ŋ	A.		
•	 10.—Compound Squills Cough	Miz	cture.
R.	Syr. scillæ	3	j

### 11.—Gastrie Catarrh Mixture.

Ŗ.	Ammonii chloridi	grs. 3	xl. ss
	Acidi hydrochlor. dil	3 3 3	j iv
1.	Syr. simplicis	3	j

### 12.—Mixture of Morcury and Jodide of Potassium.

R	Hydrarg bichloridi	gr.	ij
	Potass. iodidi	gr.	SS
	Potass. chloratis	3	
	Tr. gentianæ comp	•	•
	Aquæ	3	iv
7	M.		

# 13.—Mixture of Donovan's Solution and fodide of Potassium.

R.	Hydrarg bichloridi	grs. 3 3	í
1	Tr. cinch. comp	33	iv viij

### OF AMERICAN HOSPITALS.

## 14.—Lafayette Mixtura.

<b>R</b> .	Balsamum copaibæ.  Spt. ætheris nitrosi	333mm	j ij iv , xv vj
	15.—Lafayette Mixture (Mo:	lifie	<b>d.</b> )
	Balsamum copaibæ Liq. potassæ	CHEMENICH	j ij jss iij
	16.—Clark's Mixture.		
R	Acidi salicylici	3 3	iij ij
,	Glycerinæaa Aquæaa M.	3	ij
•			
	17.—Bismuth and Pepsin Wi	xtuı	∙•.
Ŗ	Bismuth subnitratis	3	j ss
	Glycerinæ	3	ss ij
1	M.	Ü	•

### EMULSIONS.

18.	.—Emulsion of Cod-Liver Cit phosphites.	and	Нуро-
B.	Olei morrhuæ	3	iij q. <b>s.</b> ij
19.	.—Emulsion of Cod-Liver Oil a	nd G	entian.
R.	Olei	3	ij
	0		
	PILLS.		
	20.—Arsanious Acid Pi	11.	
R.	Acidi arseniosi Cinchonidiæ sulph Pulv. capsict Ext. cinchonæ. Pulv chinoidini	gr. grs. grs.	XXX iv

	21,—Aloos and Belladonna I	PIII (F	B.).
R.		grs.	-
	Cinchoniæ sulph	grs.	
	Ferri sulph. exsic	grs.	
	Ext. belladonnæ	Rrs.	**
	Ext. nucis vomicæāā	gr.	v
	M. et ft. pil. No. xx.	g.,	•
•	M. et a. pn. 140. xx.		
22	2.—Aloos and Hyescyamus i	<b>2111</b> (1	R. H.)
R.	Pulv. aloes soc		
-	Quiniæ sulphatis		
	Ferri sulphatis exsic	grs.	xx
	Ext. nucis vomicæ	grs.	
	Ext. hyoscyami	grs.	
•	M. et ft. pil. No. xx.	•	
	23.—Strychnia and Podophyi	lum I	PIII.
R	Strychnia sulphatis	gr.	
ж	Resinse podophylli	grs.	
	Ext. colocynth. comp	grs.	
	Pulv. capsici	grs.	
	M. et ft. pil. No. xl.	B.3.	-
	POWDERS.		
	24.—Bismuth and Sod	<b>a.</b>	
R	Bismuth and subcarb		_
IX.	Sodii bicarbonatis	grs. :	
1	M. et ft. chart. No. j.	-	-

### OFFICIAL FORMULÆ

	25.—Calomei and Chaik	( <b>A.</b> )	
R.	Hydrarg. chlor. mite	gr. grs.	viij
_			
	28.—Rhubarb and ipec	BC.	
R.	Pulv. ipecacuanhæ	grs. grs.	xij
7	Pulv. cubebse	grs.	xxiv
-	u. et it. chart. 140. vj.		
	27,—Magnesia and Potass	dum.	
Ŗ	Magnesii sulphatis	3	
I	M. et ft. chart. No. j		
	28.—Santonin (C. V. C.)	)	
Ŗ	Pulv. santonini	grs.	x
	Hydrarg. chlor. mite	grs.	
	Resinæ jalapæ	gr.	
	Sacch. lactis	grs.	XX
1	M. et. ft chart. No. vj.		
	28.—Magnesia and Sulpi	hur.	
R	Magnesii		
~~	Magnesii sulphatis		
	Sulphuris		
	Sacch. albæāā	3	iv
	Pulv. anis	3	ij
	M. et. ft. chart. No. i.	_	

1/

	OF AMERICAN HOSPITALS.			187
D	30.	_		•
R.		3	)	
Ŗ.	31. Potassii chloratis (in white paper)	3	SS	
R.	32. Zinci sulphatis (in blue paper)	3	55	
	0			
	FOR EXTERNAL USE.			
	33.—Aconite Liniment.			
R	Ext. acoaiti rad. fl	NONOMO	j ss ij	
1	M.			
	34.—Rhaumatic Linimen	t.		
R.	Tr. capsici			
	Ol. olivæ	3	j	
,	Tr. origani	3	ij	

	35.—Red Wash.	
Ŗ		grs. ij
	Tr. lavandulæ comp	m xv
. 1	Aquæ	<b>3</b> j
•		
	36.—Carbelized Wash.	
R	Acidi carbolici	
	Glycerinæāā	z i
	Aquæ	g j
V	M.	
_		
_	37.—Evaperating Lotion (Conce	
Ŗ		3 jss 3 ijss
	Acidi acetici	Ž ijss
	Spt. vini rectificatiq. sad	<b>3</b> iij
I	M.—Sig. Dilute with one quart of water	•
	38.—Carbelized Oil.	
R	Acidi carbolici	7 i
1,X	Olei olivæ	3 j 3 ij
1	M.	3 3
-		
	39.	
R	Atropiæ sulph,	grs. ij
	Aquse	grs. ij Žj
	M. et ft solutio.	_
D	40.	
Ŗ	Atropiæ sulph	grs. iv
	Aquæ	<b>3</b> j
	M. et ft. solutio.	

## OINTMENTS.

	41.—Carbelic Acid Cintme	ent.	
R.	Acidi carbolici	3	j j
	42.—Carbelized Zinc Qinta	nent	<b>.</b>
R N	Acidi carbolici	3	j j
		_	
	43.—Tannic Acid Cintme	nt.	
R.	Acidi tannici	3	ss j
	44.—Boracic Acid Cintme	mt.	
Ŗ	Glycerit. acid. boracic. sat	3	ij
	Ceræ albæ	7	i
	Ung. petrolii	3	yj
b	d. et ft. unguentum.	_	

4	5.—Qintment of	Donn		Bai	sam	of
₿	Jodiformi		•••••	3 3	j ij	
	Glycerinæ	•••••		3	j	
	46.—Cbryse _l	 phanic Ac	id Oi:	atme	nt.	
B	Acidi chrysophan Unguenti petrolii M. et ft. unguentum.	ici			ij xvj	
	47.—Ca	iemol Oin	tment	ì.		
R	Hydrarg. chlor. m Ung. aquæ rosæ Ol. amygdal dulc M. et ft. unguentum.				ij ss xvj q. s	
	48.—Ammonia	tod Merc	ury Ol	ntm	ont.	
B	Hydrarg. ammonia Ungt. petrolii M. et ft. unguentum.	• • • • • • • • • • • • • • • • • • • •	•••••	3	vj xvj	
	48.—Yollow Gxi	do of Mer	cury (	Dint	mont	•
Ŗ	Hydrarg oxidi fla Ung, petrolii M. et ft. unguentum.	V	******	grs.	xvj j	

	OF AMERICAN HOSPITALS.			191
	* 50.—Diacheion Ointment—H Olei olivarum	Ð	j iij ss ij	
	Dr. Harrigan's Cough Mix	ure	<b>)</b> .	
	(Used in St. Mary's Hospital, Brookly	n, N	. Y.)	
Ŗ 1	Codeiæ	gr. M 3	X iij j	

## MEDICAL FORMULARY

OF THE

# CITY*HOSPITAI

OF THE

CITY OF BOSTON.

### SOLUTIONS.

A.

	Gramm
<ul> <li>R Zinci sulphatis</li></ul>	gr. iv   .26
A. Dilut.	<b>:</b> :
R Zinci sulphatis	gr. ij .13 <b>3</b> j 30.00

### OF AMERICAN HO

	A.—1.—
R	Zinci sulphatis
N	И.
	A1
R	Zinci sulphatis
1	И.
	Pilocarp
R	Pilocarpinæ hydrochloratis.
1	И.
	Atropin
R	Atropinæ sulphatis
1	1.
	В.
R	Sodii boratis
2	м.
	C.
R	Potassii chloratis

	v.		
R.	Acidi sulph. dil	gr. viij m. viij 3 j	.52 .52 30.00
,	$\mathbf{M} - 3 \; \mathbf{j} = \mathbf{gr. j.}$		
	F.		
Ŗ.	Ferri sulphatis	gr. iv gr. ix m. xx 3 j	.26 3.88 1.30 30.00
1	M.		
_	<b>0.</b>		
K.	Tr. ferri chloridi	m lxxx	5 00 30.00
1	$M3 j = m \times .$	3,	<b>J</b>
	P.		
Ŗ	Potass iodidi	gr. xl	2.50
3	$\mathbf{M} \cdot \mathbf{-3} \; \mathbf{j} = \mathbf{grs.} \; \mathbf{v}.$	9,1	30.00
	Р. В.		
	Potass. bromidi	gr. lxxx	5.00 30.00
1	$\mathbf{M3} \ \mathbf{j} = \mathbf{grs.} \ \mathbf{x.}$		

OF	AMERICAN	HOSPITALS.
----	----------	------------

195

### D. C.

### Chloral.

### Sodii Salicvias.

 R. Acidi salicylici...
 gr. lxv
 4.21

 Sodii bicarb...
 3 ss
 15.00

 Aquæ...
 3 j
 30.00

 M.
 3 j
 30.00

### Ferri et Quininæ Citres.

### MIXTURES.

B.	Sodii bicarb	3 ss ij o ij	Grammes. 15.0 7.5 3.8 960.0
	II.		
R.	Hydrarg. chlor. corros  Potass. iodidi	gr. 1/3 gr. xl 3 j	2.50 30.00
	III.		
	Ferri sulphatis Syrupi Tr. quassiæ	gr. xxx	1.94 15.00
B	Hydrarg. biniodidi	gr. 2–5 gr. xl m viij m cccxx 3 j	.03 2.59 0.52 20.74 30.00
:	M.		

	OF AMERICAN HOSP	ITALS.	197
R.	Morphinæ sulphatis Vini ipecac Syrupi senegæ " scilæ Aquæ	gr. ½   3 ij	.032 7 780
R.	Ammon. chlor	gr. lxxx 3 iv 3 iij 3 ij	5. 15. 10. 60.
•	Olei morrhuæ	3 ss	15.00
B.	Hydrarg chlor, corros Potass, iodidi Tr. gentian, comp Aquæ	gr. ss gr. xl ss ss	.032 2.500 15.000
Ŗ.	Liq. ammon. acet	<b>3</b> j	30.00

X.		
Tr. camphoræ Tr. opii	<b>z</b> ;	30.000
	9 ) (	30.000
<u> </u>		
XI.		
Hydrarg. chlor. corros Tr. ferri chlor	gr. j 3 ss 3 ij ss	.064 15.000 77.000
<b>1.</b>		
Spt. ætheris nit Tr. opii deodor	m x g j m xxx	.97 7.40 1·94 30.
•	37)	3
XIII.		
Morphinæ sulph	gr. ss m xx	.032 1.30
	7 55	15.00
-	3 33 1	13.00
<b></b>		
XIV.		
Morphiæ sulph Sodii "	gr. ss gr. xxiv Z j	.032 1.560 30.000
	Tr. opii Tr. rhei	Tr. opii

### OF AMERICAN HOSPITA

		XV.
R	Chloroformi	
70	Syr. pruni virg Syrupi	āā
1	M.	
		_
		XVI.
R		************
	Morphiæ sulph Spt anisi Syr. p'uni virg	
N	t.	
		-
		XX.
R	Olei lini sem	
	Syrupi Olei amygdalæ ar	āā
1	M.	
	WA	SHES, ET
	C	arbolic S
		1-80.
	Acidi carbolici	
-	M.	3

### Carbolic Sol.

### 1-40.

### Carbelic Sol.

### 1--20.

### Carbolio Oil.

### Chiorinatod Soda Wash.

### Lead Wash.

("Liq. Plumb. subac. dil., U.S.P. 1880.)

	Myrrh Wash.		
Ŗ.	Tr. myrrhæ	<b>3</b> j (	I.
	Aquæ	<b>3</b> iij	3.
1	M		
	Nitric Acid Wash		
R	Acidi nitrici		τ.
~~	Aquæad	3 j 3×vj	128.
	·	<b>J</b> , (	
	Lead and Opium Wa	ash.	
Ŗ.	Tr. opii	3 ij	I.
	Liq. plumbi subacet. dil	3 xiv	8.
	M		
	Soda and Opium Wa	-h	
R	-		_
IX.	Liq. sodæ chlor Tr. opii	rongra j x	2. T.
	Aquæ	3 x	10.
7	M. ·	•	
	Chleroform and Aconite I		
Ŗ		3 j	30.
	Chloroformi	3 j	
	и.	3 35 1	15.
•			
	Evaporating Lotio	n.	
R	Ammonii chlor	3 i l	ı.
_,-	Acidi acetici dil	3 j	_
	Spt. vini rect	<b>-</b> .	
1	Aquæ	3 W	3 <b>2</b> .

	Mouth Wash.		
Ŗ	Glycerini	3 ss	
_	Aquæ rosæāā	3 ss (	15
I	М.		
		•	
	. Alkaline Powdo	rs.	
Ŗ.	Potassii bicarb	1	
	Sodii	gr. xv gr. ij	.97
_	Pulv. zingiberis	gr. ıj	.13
1	M.		
	OINTMENTS		
	OINTMENTS	•	
		G	rammes
Ŗ.	Ung. hydrarg. nit	3 j (	I.
	Ceratum	3 j 3 iij	3.
2	М.		
ĸ	Pulv. iodiformi Ung. petrolei	gr. xiviij	10. 90.
		3) (	90.
	м.		
	PILLS.		

.01 .13

OF AMERICAN HOSP	ITALS.	203
R Ferri sulph. exsic Ext. nucis vom	gr. j gr. ¼ gr. j	.065 .016 .065
### Ferri sulph. exsic	gr. 3-5 gr. j	.039 .065
R. Pulv. camphoræ	gr. j	.065
R. Ext. colocynth. comp	gr. iij   gr. 1-6	.194
VI.           Pulv. capsici	gr. ij gr. ss gr. j	.130 .032- .065

M.

	VII.		
Ŗ	Pulv. opii cupri sulph	gr. j   gr. ¼	.065
	<b>M.</b>		
	VIII.		
Ŗ.	Pulv. digitalis	gr. j gr. ss gr. 1-6	.065 .032
	₫.		
_	ıx.	•	
Ŗ	Ferri redactiQuininæ sulphāā Strychninæ sulph	gr. j gr. 1-60	.065
	<b>V</b> .	•	
	X.		
	(Pil. Ferri comp's., U.S.P	., 1880.)	
Ŗ	Pulv. myrrhæ	gr. j as	.097
	Pulv. myrrhæ Sodii carb Ferri sulphāā	gr. 34	.09 <b>7</b> .048
	<b>M.</b>	- ,,	
	XI.		
Ŗ	Ext. hyoscyami	gr. ij	. 1 30

		205
XII.	•	
R Pulv. digitalis	gr. ss gr. j gr. j ss	.032 .065 .097
M.		
XIII.		
Res. podophyllin	gr. ½ gr. iij gr. j	.022 .195 .065

## CHICAGO MARINE HOSPITAL.

U. S. MARINE-HOSPITAL SERVICE,

(District of the Great Lakes.)

PORT OF CHICAGO, ILL., SURGEONS'S OFFICE, MAY 15, 1885.

Dr. C. F. TAYLOR,

Philadelphia.

Sir:—I have the honor to herewith transmit the formulæ in use at this hospital. Nos. I to 60 are kept constantly on hand, though the practice is not routine in the hospital here, and the R's are prescribed where they meet the indications. They are not all original, and I would give credit if I knew to whom credit was due. I have no doubt you will find many of them duplicated from other sources. Where a "break" in numbers occurs it is by reason of retaining the hospital number, and well-known prescriptions like "Squibb's Mixture," "Brown-Sequard's Pills," etc., have been omitted. I am not responsible for the miserable Latin in some of the formulæ, as I could not write them myself.

Yours truly,
W. H. Long, Surg. M. H. S.

(Note.-30 cubic centimeters equal 3 1.01. For

practical purposes, to reduce cubic centimeters to ounces, divide the number of cubic centimeters by thirty. One gram equals 15.432 grains, or about 15½ grains. To reduce grams to grains, multiply the number of grams by 15½.)

### 1.—Fever Mixture.

R	Tr. aconit. rad	20	c. <b>c.</b>
	Spt æther nit	200	44
	Tr. cardam. comp	150	"
	Liq. ammon. acetad	1000	66
1	M.—Dose, five cubic centimeters.		

### 2.-Fever Mixture.

Ŗ.	Tr. aconit. rad	30	
•	Tr. gelsemii	100	"
	Tr. opii camph	150	"
	Pilocarpin mur	1	50 gm,
	Liq. ammon. acetad	1000	c. c.
7	L.—Dose, five c. c.		

### 3.—Fever Mixture.

Ŗ	Acid. hydrobrom. dil	300	c. c.
	Quiniæ sulph	20	gm. c. c.
	Spt. chloroformiad	300 1000	

M.—Dose, five c. c.

#### 4.—Rheumatic Mixture.

Ŗ	Potass. iodid	150	gm. c. c.
	Tr. cimicifugæ	300	"
	Syr. sarsap. comp	300	"
	Aquæad	1000	**
1	M.—Dose, five c. c.		
	5.—Rheumatiç Mixtur	<b>e.</b>	
Ŗ	Acid. salicyliciPotass. bicarb	50 100	gm.
	Tr accrit rad	- 1	

#### 

Aquæ.....ad... 1000

M.—Dose, five c. c.

#### 6.—Syphilitic Mixture.

$\mathbf{R}$	Hydrarg. chlorid. corros	1	gm.
	Potass. iodid	100	٠.,
	Tr. gentian. comp	300	c. c
	Extr. sarsap. comp. fl	100	66
	Aquæad	1000	"
	Aquæad	1000	

M. -Dose, five c. c.

#### 7.—Syphilitic Mixture.

Ŗ	Potass. iodid Extr. stilling. fl	200	gm. c. c.
	Tr. cinchon. comp	300	"
	Aquæad	1000	"

M.—Dose, five c. c.

	8.—Cough Mixture.		
Ŗ	Ammon. chlorid	100	gm.
	Syr. senegæ	150	c. c.
	Syr. scillæ	150	
	Syr. pruni virg	300	46
	Morphiæ chlorid	1	gm.
	Extr. glycyrrhizæ fl	100	c. c.
	Aquæad	1000	46
	M.—Dose, five c. c.		
	10.—Cough Mixture	•	
Ŗ	Ammon. chlorid	100	gm.
	Potass. bromid	100	16
	Tr. opii camph	200	c. c.
	Syr. ipecac	200	"
	Syr. tolutan	300	"
	Ext. sarsap. comp. flad	1000	46
ļ	M.—Dose, five c. c.		
	11.—Rheumatic Mixtu	re.	
R	Potass. iodid	100	gm.
	Ammon. chlorid	ICO	٠,,
	Tr. digitalis	100	c. c.
	Pilocarpin. chlorid	2	gm.
	Mist. glycyrrhizæad	1000	č. c.
]	M.—Dose, five c. c.		
	13.—Diarrhœa Mixtur	θ.	
Ŗ.	Tr. opii camph	200	c. c.
	Tr. catechu	200	46
	Tr. cardam comp	100	**
	Mist. cretæad	1000	"
	M.—Dose, five c. c.		

#### 14.—Tenic Mixture.

Ŗ.	PepsinæQuiniæ sulph	20	gm.
	Acid. nit. muriat. dil	50	<b>c</b> . c.
	Glycerinæ	100	"
	Aquæad	1000	"
1	M.—Dose, twenty c. c.		

#### 15.-Laxative Mixture.

R	Magnes. sulph	250	gm.
	Ext. glycyrrhizæ	100	٠,,
	Ext. sennæ fl	50	c. c.
	Elix. aurantii	250	"
	Aquæ bullientad	1000	"
1	M.—Dose, one hundred c. c.		

#### 18.—Diapheretic and Diuretic.

Ð	Tr. digitalis	•	٠
ĽX.	Tr. digitalis	100	
	Potass. iodid	100	gm.
	Potass acet	200	"
	Pilocarpin. chlorid	1	.50 ''
	Spt. æther nit	300	c. c.
	Syr. scillæ	150	"
	Syr. simplicisad	1000	**

M.—Dose, five c. c.

## OF AMERICAN HOSPITALS.

	17.—Diuretic.
R	Potass. bicarb
**	Ext, buchu fl
	Ext. uvæ ursi fl
	Tr hyoscyami
	Acaciæ pulv
	Olei sassaf
	Aquæ camphad
1	M.—Dose, five c. c.
	18.—Gonorrheal Mixtu
B	Copaibæ bals
	Liq. potassæ
	Spt. æther nit
	Spt. lavand. comp
	Syr. simplicisad
1	M.—Dose, five c. c.
	The second second
	19.—Gonorrheal Mixtu
R	Gurjun bals
**	Liq. potassæ
	Tr. cardam comp
9 4	Syr. simplicisad
1	M.—Dose, five c. c.
	The state of the s
	20.—Tonic.
13	
B	Tr. ferri chlorid
	Tr. cantharidis
	Elix aurantii
	Aquæad
1	M.—Dose, five c. c.
-	The state of the s

#### OFFICIAL FORMULÆ

#### 21.

R.	Zinci. sulph	I I I I I I I I	gm. .30 " .50 " c. c.
	22.		
Ŗ.	Acid. carbol. cryst	20 4 50 2 1000	gm. " c. e. "
	<del></del>		
	23.		
Ŗ.	Ext. hydrast. fl	100 100 250 1000	c. c. " "
	25.—Tonic.		
B.	Ferri et potass. tart	200 200 1000	gm. .30 " c. c. "

#### OF AMERICAN HOSPIT

	201 onic,
R	Quiniæ sulph
1,8	Tr. ferri chlorid
	Spt. chloroformi
	Glycerinæ
	Aquæa
	The second secon
-	M.—Dose, five c. c.
	27.—Tonic.
R	
IX	Cinchonæ sulph Tr. ferri chlorid
	Spt. chloroformi
	Elix, aurantii
	Aquæa
3	M.—Dose, c. c.
	8.—Anti-Perio
R	
UK.	Quiniæ sulph
	Acid. sulph. arom
	Tr. capsici
	Syr. rhei. arom
3	M.—Dose, five c. c.
	30.—Tonic.
51	Acid. nit, muriat. dil
107	Tr. calumbæ
	Tr. cardam. comp
	Civetine
	Glycerinæ
-	Aquæ

## 31.

B.	Chloral hydrat	200 200 200 300 10	gm. « c. c. « «
	<b>32</b> .		
B.	Chloroformi	300 50 200 300 1000	c. c. " "
	33.		
R	Chloroformi	50 50 50 500 1000	c. c.
	VI.—External use.		
	34.		
<b>₽</b> .	Peruviani bals Tr. myrrhæ Benzoini compad M.—External use.	300 300 1000	c. c. "

#### 35.

R N	Ammoniæ chlorid	300 1000	gm. c. c.
	36.		
Ŗ,	Tr arnicæ	250 250 1000	c. c. "
	<del></del> 37.		
Ŗ.	Iodini	15	gm. c. <b>c</b> .
	-		
R N	Potass. chlorat	50 30 250 1000	gm. " c. c. "
	 39.		
B.	Acid. tannici	30   250   1000	gm. c. c.

	ı.

R.	Acid. boracici	150   850	gm
	Acid. carbol. cryst	50   950	gm "
R M	Ung. hydrarg. nit	100 100 100 700	gm. "
	Ung. hydrarg. nit	200 800	gm.
R.	44.  Iodiformi	20 100 880	gm.

#### OF AMERICAN HOSPITALS.

		45.
R	Zinci oxidi	
N	I.—External use.	
		_
		46.
R	Hydrarg, ammoniat Tr. benzoin. comp Ung. petrolei	*******
N	1.—External use.	and III-li
		47.
	Acid, crysophanici Ung. petrolei M.—External use.	
		48.
R	Bismuthi subnit Sodii bicarb	
1	M.—Dose, one gram.	
		49.
B	Magnesii sulph	************
	Sulphuris Glycyrrhizæ pulv	
	M.—Dose, one gram.	

500	gm.
500 500	gm.
	<b>5(</b>

#### 52.—Tertiary Syphilis.

Ŗ.	Iodinii resublim	4   80	gm.
	Potass. iodid	8o	"
	Syr. sarsap. compad	1000	c. <b>c</b> .
1	M —Dose five c c		

#### 53.—Tenic.

R	Quiniæ sulph	13	gm.
~~	Tr. ferri. chlorid	66	č. c.
	Strichniæ sulph		20 gm.
	Aquæad	1000	c. c.
7	/Dose, five c. c.		-

#### 64.—Syphilitic Mixture.

Ŗ	Potass. iodidAmmon muriat	120	gm.
	Aquæad	1000	c. c.
N	1.—Dose, five c. c.		

I) gm.

30 gm.

30

30

120

60

30

1000

c. c.

66

66

#### 55.—Syphilitic and Tonic.

Hydrarg. bichlorid.....

Ammon sesq. chlorid	1	~"
Quiniæ sulph		46
Aquæad	1000	"
M —Dose, five c. c.		
56.		
R Zinci sulph	61	øm.
Morphiæ sulph	2	5
k Zinci sulph	1000	c. c.
M.—Injection.	1000,	
58.—Tonic.		
R Tr. ferri chlorid	20	c. c.
Liq. potass. arsen	12	c. c. "
Quiniæ sulph	4	gm.
M.—Dose, one c. c.	. "	•
59.—Cough Mixture	) <b>-</b>	

Glycyrrhizæ pulv.....

Sacchari alb.....

Acacise pulv.....

Tr. opii camph.....

Vin. antimonii.....

#### 

R

#### 60,-Catarrh Snuff.

R	Bismuthi subnit	8	gm.
	Sacchari alb. pulv	16	"
	Camphorse pulv	2	03 C. C
1	Olei gaultheriæ	U	03 0.0

## COOK COUNTY HOSPITAL,

(CHICAGO, ILL.)

W. J. McGarigle-Warden.

CHICAGO, April 20, 1885.

#### DR. C. F. TAYLOR.

R

Yours at hand. We have but few established formulæ in use here, the hospital having assumed such dimensions as to demand a drug room in each ward. The following are those in use at present:

#### Mist. Ammon. Murias.

Ŗ.	Ammon. mur Morphiæ sulph	grs.	
	Antimon, et pot. tart	grs.	
N	1.	0	~',

#### Hydrocyanic Acid Cough Mixture.

Acidi hydrocyanici dil	3	j
Syr. ipecac	3	j
Syr. tolutaniad	3	viij

#### Liniment.

R Aquæ ammon	Z iv
Tr. arnicæāū.	Z iv
Aquæ	•
M.	

#### Aiterativo.

R	Iod. potass	3 viij
	Syr. sarsap	Z viij
M	4. ·	-

Very respectfully yours,

F. P. MURPHY, Druggist.

# MASSACHUSETTS GENERAL HOSPITAL.

(BOSTON.)

Boston, March 3, 1885.

DR. C. F. TAYLOR.

Dear Sir:—In reply to yours of the 24th inst. would say, we have no printed formulary, but have a number of prescriptions which are peculiar to the Massachusetts General Hospital, and only used by the hospital staff. I will give you the R's that I consider peculiar to the hospital, and if I can give you any further information would be very glad to do so.

Yours very truly,

Wm. L. A. CAMERON,

For the Sup't.

Apothecary.

#### Hospitai Pilis.

R	Ex. colocynth comp	grs. iij gr. j
	Nucis vomicæ	gr. 1
М	Ft. pil, No. j.	

	Hospital Diarrhœa Mixture.
₿	Tr. cinnamomi Tr. opii
	Tr. catechu
	M.—Dose, 3 ss.
	Hospital Solution Saits.
R	
~	Aq. menth. pip
	Aquæ
	M.—Dose, 3 iv.
	Hospital Cough Mixture.
R	
	Morph. sulph gr. j Syr. simp 3 iv
	Chloroformi n sv
	M.—Dose, 3 j.
	Hespitai Fovor Mixture.
R	Spt. æth. nit
	Spt. chloroformi
	Liq. ammon acet
	M.—Dose, 3 j every hour.
	Shaw's Specific.
R	•
	Hydrarg. biniod gr. 1-12
	Aquæ
	Syr. zingib 3 ss
	M.—Dose, 3 j, three times daily.
	, O 1,

## MARYLAND WOMAN'S HO

(BALTIMORE, MD.)

#### House Tonic.

R	Tr. ferri. chlor	
	Strychinæ sulph.	***************************************
	Aquæ	
1	MFilter, Sig.	One teaspoonful the

M,-Filter. Sig. One teaspoonful threafter meals.

#### House Injection.

R	Zinci sulph
	Acidi carbolici
	Aquæ

M.—One ounce to a pint of water. U injection; from one to three syringefuls fr times a day.

These are the only established formul is treated according to indications present

## ST. JOHN'S HOSPITAL.

(BROOKLYN, N. Y.)

St. John's Hospital, Brooklyn, N. Y., April 27, 1885.

Dr. Taylor,

Dear Doctor:—You was here a short time since and wanted me to give you a few prescriptions which we have used with advantage in the hospital. The following are such:

#### Goneral Sedative Cough Mixture.

R.	Morphiæ sulph	grs.	iv
	Acidi hydrocyanici diluti	fl. z	ij
	Acidi hydrocyanici diluti Syrupi tolutan	fl. 🕉	viij
7	A —Sig Teaspoonful as needed	_	_

#### Diarrhosa Mixture.

R.	Tr. catechu			
	Tr. opii camphāā	fl	3	i
	Tr. opii <b>c</b> amph	fl	3	ij
N	Sig One or two teaspoonfuls as ne	ede	ď.	-

#### In Cystitis.

R	Fl. ext.	. bucl	nu comp	 				
			•••••		f	3	÷	s;
1			teaspoonful					

#### Another.

R	Acidi benzoici	grs. lxxx
	Sodæ bi-boras	3 ij
	Aquæ menth. pip	
:	1.—Sig. Tablespoonful three time: 2 d	ay.

#### A Tonic Pill to be Used in Chronic Constipation.

Ŗ.	Ext. nucis vom	āā	grs. v grs. x
	Ferri sulph. exsic	āā	•
M day	f.—Make into twenty pills.	Sig. On	e three times a

I may send you more at another time.

Very respectfullv,

Resident Physician.

#### ST. LOUIS CITY HOSPITAL.

(ST. LOUIS, MC.)

OFFICE OF SUPERINTENDENT OF CITY HOSPITAL, St. Louis, March 22, 1885.

C. F. TAYLOR, M. D., Philadelphia, Pa.

Dear Sir .- Your letter of the 18th inst. received. We have no printed formulary, and our established formulary is only for antiseptics after Lister, Nussbaum, etc... which, of course, everyone has. We prepare our own gauze from caleramic cloth, using, however, iodoform dry dressings for wounds, abscesses and the like. I use five per cent, carbolized boiled linseed oil for burns and frost-bites, leaving them exposed to the air without further dressings. We use boric acid solutions in washing cavities after aspirating, where we do not have to resort to free opening. In the latter resort, we commonly employ two and a half or five per cent, solutions of carbolic acid Corrosive sublimate solution, one part to a thousand, we use on occasion. I use the metric system in prescribing-have done so for six years-and every dose given to our nearly seven thousand patients a year is prescribed especially; i. e., a small bottle or quantity at a time. We have no standard mixtures or prescriptions, but vary according to indications. Cinchonidine has been used exclusively instead of quinine for the past nine years with the best results, and not in large doses at that. To mention some particular way more often used in giving a particular remedy, like equal parts of salicylic acid and acetate of potash in syrup of lemon, is too elementary and didactic.

Yours truly, D. V. DEAN.

Supt. and Surgeon in charge.

## CINCINNATI HOSPITAL.

(CINCINNATI, OHIO.)

CINCINNATI, O., March 20, 1885.

Dr. C. F. Taylor,

Editor Medical World.

Dear Sir:—Yours of the 18th, addressed to the superintendent of this hospital, has been referred by him to me, for answer. The Cincinnati Hospital has no printed formulary, and does not use one.

Our list of drugs and chemicals include everything

that is used by the regular schools of medicine.

The hospital is conducted on a very liberal plan, and all new remedies are given a fair and thorough test as soon as they are obtainable; and, if found really valuable, their use is continued. Many remedies are in constant use in the hospital which are as yet comparatively unknown outside.

Very few actual prescriptions, as that word is commonly understood, are written. There is kept in every ward a large, portable medicine case which contains all the remedies in general use in that ward, the medicines being kept in the form of solutions, powders or pills, most convenient for administration. The resident physi-

cians in making their rounds prescribe simply the required number of grains, or drachms, etc., of the medicine, and they are obtained from this stock.

New prescriptions are only written when some new remedy is desired, or certain combinations which cannot be made up from the ward supply. The ward supply is replenished daily from the druggist's department, and is frequently overhauled by him.

In the same way every ward is supplied with all the instruments in ordinary use in that ward.

The subject of treatment is a very large one, and I take it, you want only a general answer. The hospital staff, as a rule, believe in very little medication.

In the treatment of continued fevers, we make large use of external cold, either in the form of sponging, ice caps, cold baths. the wet pack, or ice-water coils.

Quinia is used extensively, but most of the stronger preparations, as kairin, are no longer used, being considered too depressing and dangerous.

In most diseases the principal object in the treatment is to build up and sustain the strength of the patient until the affection has run its course.

The surgeons use very extensively iodoform, powdered boracic acid, mercury bichloride solution, and in less amount, carbolic acid.

Dr. E. W. Walker is probably entitled to priority in the use of boracic acid powder for the treatment of compound fractures.

Fractures are dressed almost entirely by plaster of Paris applications, either modified Bavarian, or made of plaster rollers.

For fracture of the femur, Dr. D. S. Young has a very excellent, yet simple device of his own, which makes

extension by the use of weights, and supports the entire limb in a shallow box, so that the thigh can be examined in its entirety, without any movement, or pain.

in its entirety, without any movement or pain.

The anæsthetics used are ether and the a. c. e. mixture. The muriate of cocaine is used as a local anæsthetic by the oculists and gynæcologists.

Yours etc.,

FRANK W. HENDLEY, M. D., Senior Resident Physician, Cincinnati Hospital.

# WESTERN PENNSYLVANIA HOSPITAL FOR THE INSANE.

(DIXMONT, PA.)

DIXMONT, ALLEGHANY Co., PA., March 21, 1885. Dr. C. F. TAYLOR:

DEAR DOCTOR: In answer to your letter I may say that we have no regular form of medicines here, in fact we give very little. Many of our patients are broken down physically as well as mentally, and we depend on tonics, etc., to build them up and return them to health of body and mind.

We use almost no sleeping medicine, no chloral nor hyoscyamin, and only lupulin when really necessary, and little or no restraint.

Yours very truly,

H. A. HUTCHINSON,

Medical Superintendent.

## U. S. NAVAL HOSPITAL.

(WASHINGTON, D. C.)

U. S. NAVAL HOSPITAL, WASHINGTON, D. C., April 28, 1885. C. F. TAYLOR, M. D.,

Editor Medical World,

Dear Doctor:—I was in Philadelphia when yours, of 18th, arrived, and regret not having seen you to explain in person that we have no printed formulary for use in the medical corps, and that I discourage set prescriptions for routine treatment in this hospital. Careful nursing and sanitation constitute a very great part of our Naval Hospital treatment.


The sick sailor, after a warm bath, clean clothes, and rest in a comfortable bed, often requires but little else. We feed the men well and dose them as little as possible. chronic pulmonary phthisis and venereal affections constitute the larger number of our patients, and they receive the customary medical treatment.

Faithfully yours,
ALBERT L. GIHON,
Medical Director, U. S. N., in charge.

## U. S. MARINE HOSPII

(SAN FRANCISCO, CAL.)

U. S. MARINE HOSPI SAN FRANCISCO, Cal., May



### HOME OF INEBRIATES.

(SAN FRANCISCO, CAL.)

SAN FRANCISCO, Cal., March 20, 1885.

C. F. TAYLOR, M. D.

My Dear Sir:—In reply to yours of the 13th inst., I have to say, we have no printed formulary in this hospital. Each case is treated separately and distinctly, and entirely on its merits. We do not use alcohol in any form. We do use milk, the very best, Jersey dairy, extensively, in treatment of alcoholic cases.

To procure sleep, we use bromide potassium and chloral hydrate.

B.	Potass.	bromidi	
	Chloral	hydratisāā	3 j
		••••	₹ j U j

M.—Sig. Of this mixture, one to two ounces every four hours until sleep is procured. If one ounce is enough, well and good; but we use two ounces if necessity require.

This does not often fail us, but when it does, we combine with each dose a drachm of tinct. hyoscyamus; or we give a hypodermic injection of morphia, say gr. ¼, and repeat in two hours if necessary. The principal

thing is to overcome nervous action, and procure sleep. Immediately afterwards, we produce evacuations from the bowels and look well to the bladder. To support the patient we use milk, beef tea and soup; and to stimulate, we use acetate of ammonia in full doses. Each patient is seen twice daily, at 9 A. M. and 6 P. M., and as much oftener as necessary. If he shows a tendency to sink, the battery is used freely and long continued.

The food is of the most nourishing character. An excellent library, cards, chess, checkers, backgammon, dominoes and pictures in great profusion are used to take the patient's mind off his disease, and with very good effect.

Yours Truly,

J. G. JEWELL,

Resident Physician and Superintendent.

N. E. cor. Stockton and Chestnut streets.

SAN FRANCISCO, Cal., March 20, 1885.

C. F. TAYLOR, M. D.,

Editor Medical World,

Philadelphia, Pa.

Dear Sir:—I received yours of the 13th inst. and also the copy of the chart "The Urine in Disease," for which please accept my thanks.

Dr. Alers and myself have given up our Sanitarium for Alcoholism and Opium Habit, as we found it would not answer without some *law* enforcing *restraint*, which is wanting in this State. While under my charge (as resident physician) the formulæ used were the usual combinations of chloral hydrate and potassium bromide, given

in aquæ menthæ. In some desperate cases of delirium tremens the heroic doses of tincture of digitalis (3 ss) were tried with success.

The opium cases (as a general rule) were treated by a gradual reduction of the accustomed drug, combined with tonics and nerve stimulants. I treated successfully one case in which one drachm of morphia sulph. was taken daily. This is the largest quantity, that has actually come unite my personal observation.

I am sorry that I cannot give you any more information on these interesting topics.

With many thanks for your kind letter, I remain,
Yours truly,
A. P. HAYNE, M. D.

533 Kearney St.

THE MEDICAL WORLD has grow favor with the Profession more rapidly perother Medical Journal ever published. It is now reaches every section in the United and several other foreign countries. Its limit in the reach of all—indeed, subscriberation write to us, suggesting that we raise the period with the property with the property of th

All communications should be addressed

THE MEDICAL WO

# THE CHART "THE URIN

to commended by seem and Medical Press.

1 . . . tor each abnormal ingredient, 1 y control quantity, characters, ren notes, one care shown at a glance.

· Doggibart is an invaluable exhibit to or the essential points in urinology." - Z

Physica handy chart to have in the c the got more solid information from a glan you can in a half hour pouring over text 1 Sugar Sweet Sword of Medicine


Worth double the money." - Therapeutic

by I to browler, of Cochton, N. Y., wri woulder how much function knowledge can be couls according and confensed in so small a sp lilitique

THE MEDICAL WORLE 1820 Chroinul St. Philadel

١.







#### LANE MEDICAL LIBRARY

To avoid fine, this book should be returned

on or before the date last stamped below.				
r				
	·	1		
•				

